

Inequalities for the perimeter of an ellipse

G.J.O. Jameson, *Math. Gazette* **98** (2014)

The perimeter of the ellipse $x^2/a^2 + y^2/b^2 = 1$ is $4J(a, b)$, where $J(a, b)$ is the “elliptic integral”

$$J(a, b) = \int_0^{\pi/2} (a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2} d\theta. \quad (1)$$

This integral is interesting in its own right, quite apart from its application to the ellipse. It is often considered together with the companion integral

$$I(a, b) = \int_0^{\pi/2} \frac{1}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2}} d\theta. \quad (2)$$

Of course, we may as well assume that a and b are non-negative. We record first some elementary facts about these integrals:

- (E1) $J(a, a) = \int_0^{\pi/2} a d\theta = \frac{\pi a}{2}$, $I(a, a) = \frac{\pi}{2a}$;
- (E2) $J(a, 0) = \int_0^{\pi/2} a \cos \theta d\theta = a$, $I(a, 0)$ is undefined;
- (E3) for $c > 0$, $J(ca, cb) = cJ(a, b)$ and $I(ca, cb) = \frac{1}{c}I(a, b)$;
- (E4) $J(b, a) = J(a, b)$ and $I(b, a) = I(a, b)$ (substitute $\theta = \frac{\pi}{2} - \phi$);
- (E5) $J(a, b)$ increases with a and with b , and $I(a, b)$ decreases.

In general, neither integral is amenable to evaluation by simply writing down an anti-derivative. However, both can be evaluated in terms of the *arithmetic-geometric mean* $M(a, b)$ of a and b . This is the common limit of the sequences (a_n) and (b_n) defined by the iteration $a_0 = a$, $b_0 = b$ and

$$a_{n+1} = \frac{1}{2}(a_n + b_n), \quad b_{n+1} = (a_n b_n)^{1/2}.$$

Write $a_n^2 - b_n^2 = c_n^2$ and $S = \sum_{n=0}^{\infty} 2^{n-1} c_n^2$. It turns out that

$$I(a, b) = \frac{\pi}{2M(a, b)}, \quad (3)$$

$$J(a, b) = (a^2 - S)I(a, b). \quad (4)$$

Moreover, the iteration converges very rapidly, so these identities amount to an effective way to calculate the integrals. Identity (3) is a famous theorem of Gauss, dating from 1799. Identity (4) is the basis of the very efficient Gauss-Brent-Salamin algorithm for calculation of π . We will not enter into any details here; for a highly readable account, see [Lo].

While (4), in principle, is an exact evaluation of $J(a, b)$, it certainly does not convey a quick and transparent indication of its magnitude in terms of a and b . This purpose is better achieved by inequalities comparing $J(a, b)$ with simple expressions like $a + b$ and $(a^2 + b^2)^{1/2}$. One lower bound, $\frac{\pi}{4}(a + b)$, was given in [LS], as a solution to a *Gazette* problem. However, bounds of this sort are accurate for some values of b/a and less accurate for others, so for a really satisfactory estimation of $J(a, b)$, more than one lower bound (and equally, more than one upper bound) is needed. Here we will describe three different ways to derive such bounds directly from (1). I hope that some readers will share my view that the methods themselves are as interesting as the conclusions.

First, we mention a pair of bounds that follow at once from elementary facts (E1) and (E5): if $a \geq b$, then $J(b, b) \leq J(a, b) \leq J(a, a)$, so that

$$\frac{\pi}{2}b \leq J(a, b) \leq \frac{\pi}{2}a. \quad (5)$$

Geometrically, this is saying that the perimeter of the ellipse lies between those of the inscribed and circumscribed circles.

Another pair of bounds that seem geometrically obvious is

$$(a^2 + b^2)^{1/2} \leq J(a, b) \leq a + b, \quad (6)$$

since this says that the curve of the quarter-ellipse is longer than the straight-line path between the same points, but shorter than two sides of the rectangle. However, it is instructive to see how these inequalities can be proved analytically. For the right-hand inequality, this is very easy: since $(x^2 + y^2)^{1/2} \leq x + y$ for positive x, y , we have $(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2} \leq a \cos \theta + b \sin \theta$ for each θ . Integration on $[0, \pi/2]$ gives $J(a, b) \leq a + b$.

We will improve this inequality below, as well as giving an analytic proof of the left-hand inequality in (6). We will also establish inequalities in the reverse direction to both sides of (6), with suitable constants inserted.

Our first method uses the *Cauchy-Schwarz inequality*, which states: for non-negative numbers a_r, b_r ($1 \leq r \leq n$), we have

$$\sum_{r=1}^n a_r b_r \leq \left(\sum_{r=1}^n a_r^2 \right)^{1/2} \left(\sum_{r=1}^n b_r^2 \right)^{1/2}.$$

This can be stated very concisely in vector notation: it says $\mathbf{a} \cdot \mathbf{b} \leq |\mathbf{a}| \cdot |\mathbf{b}|$, where the length of the n -dimensional vector $\mathbf{a} = (a_1, a_2, \dots, a_n)$ is $|\mathbf{a}| = (\sum_{r=1}^n a_r^2)^{1/2}$, and $\mathbf{a} \cdot \mathbf{b}$ means $\sum_{r=1}^n a_r b_r$. The proof is short and neat, so we repeat it: writing $\mathbf{a} \cdot \mathbf{b} = S$, we have, for any λ ,

$$0 \leq \sum_{r=1}^n (a_r - \lambda b_r)^2 = |\mathbf{a}|^2 - 2\lambda S + \lambda^2 |\mathbf{b}|^2.$$

With λ chosen to be $S/|\mathbf{b}|^2$, this says $|\mathbf{a}|^2 - S^2/|\mathbf{b}|^2 \geq 0$, hence $S \leq |\mathbf{a}| \cdot |\mathbf{b}|$.

There is a corresponding statement for integrals, proved in an analogous way. We write $\int_a^b f$ (this is perfectly adequate notation!) for $\int_a^b f(x) dx$. The inequality states: *if f and g are non-negative, integrable functions on $[a, b]$, then*

$$\int_a^b fg \leq \left(\int_a^b f^2 \right)^{1/2} \left(\int_a^b g^2 \right)^{1/2}.$$

The Cauchy-Schwarz inequality enables us to give a rather elegant analytic proof of the left-hand inequality in (6), as follows:

PROPOSITION 1. *We have*

$$J(a, b) \geq (a^2 + b^2)^{1/2}. \quad (7)$$

Proof. By the Cauchy-Schwarz inequality,

$$a^2 \cos \theta + b^2 \sin \theta = a(a \cos \theta) + b(b \sin \theta) \leq (a^2 + b^2)^{1/2} (a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2}.$$

Integrating on $[0, \pi/2]$, we obtain

$$a^2 + b^2 \leq (a^2 + b^2)^{1/2} J(a, b),$$

hence (7). □

Note that (7) is exact when $b = 0$, since $J(a, 0) = a$, but not when $b = a$. A variation of the reasoning gives a second lower bound which is exact when $b = a$ but not when $b = 0$, and which also improves upon the one in (5). This bound was established in [LS], by essentially the same method (though without explicit mention of the Cauchy-Schwarz inequality).

PROPOSITION 2. *We have*

$$J(a, b) \geq \frac{\pi}{4}(a + b). \quad (8)$$

Equality holds when $a = b$.

Proof. By the Cauchy-Schwarz inequality,

$$\begin{aligned} a \cos^2 \theta + b \sin^2 \theta &= (a \cos \theta) \cos \theta + (b \sin \theta) \sin \theta \\ &\leq (a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2} (\cos^2 \theta + \sin^2 \theta)^{1/2} \\ &= (a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2}. \end{aligned}$$

Integration gives (8), since $\int_0^{\pi/2} \cos^2 \theta d\theta = \int_0^{\pi/2} \sin^2 \theta d\theta = \frac{\pi}{4}$. Equality holds when $a = b$, by (E1). □

Note that this says that the perimeter of the whole ellipse is at least $\pi(a + b)$. By the inequality of the means, this is not less than $2\pi(ab)^{1/2}$, still with equality occurring when $a = b$. As observed in [LS], this means that among ellipses with a given area, the one with the smallest perimeter is the circle. Equally, among ellipses with a given perimeter, the circle is the one with the largest area.

An upper bound for $J(a, b)$ is provided by the Cauchy-Schwarz inequality for integrals:

PROPOSITION 3. *We have*

$$J(a, b) \leq \frac{\pi}{2\sqrt{2}}(a^2 + b^2)^{1/2}. \quad (9)$$

Equality holds when $a = b$.

Proof. Apply the Cauchy-Schwarz inequality for integrals with f written as $f.1$ and both sides squared. We obtain

$$J(a, b)^2 \leq \left(\int_0^{\pi/2} 1 \right) \int_0^{\pi/2} (a^2 \cos^2 \theta + b^2 \sin^2 \theta) d\theta = \frac{\pi}{2} \frac{\pi}{4} (a^2 + b^2). \quad \square$$

Since $\pi/(2\sqrt{2}) \approx 1.1107$, the pair of bounds in (7) and (9) shows that $J(a, b)$ is actually modelled fairly well by $(a^2 + b^2)^{1/2}$.

We now describe a second strategy, based on the triangle inequality for vectors. Note that, in the notation for the length of vectors,

$$J(a, b) = \int_0^{\pi/2} |(a \cos \theta, b \sin \theta)| d\theta.$$

The familiar triangle inequality for vectors says that $|\mathbf{u}_1 + \mathbf{u}_2| \leq |\mathbf{u}_1| + |\mathbf{u}_2|$. This is geometrically obvious in the case $n = 2$, and in general it follows easily from the Cauchy-Schwarz inequality. We show that $J(a, b)$ also satisfies the triangle inequality, in the following sense:

PROPOSITION 4. *Let $\mathbf{u}_j = (a_j, b_j)$ ($j = 1, 2$), where $a_j, b_j \geq 0$. Then*

$$J(\mathbf{u}_1 + \mathbf{u}_2) \leq J(\mathbf{u}_1) + J(\mathbf{u}_2). \quad (10)$$

Proof. By the ordinary triangle inequality, we have for each θ

$$|[(a_1 + a_2) \cos \theta, (b_1 + b_2) \sin \theta]| \leq |(a_1 \cos \theta, b_1 \sin \theta)| + |(a_2 \cos \theta, b_2 \sin \theta)|.$$

Integrating on $[0, \frac{\pi}{2}]$, we deduce that

$$J(a_1 + a_2, b_1 + b_2) \leq J(a_1, b_1) + J(a_2, b_2),$$

which equates to (10). □

Note. Since $J(a, b) = J(|a|, |b|)$, (10) actually holds without the condition that the components are non-negative. In the usual parlance, $J(a, b)$ is a *norm* on the vector space \mathbb{R}^2 , and we are comparing it with other norms like $|a| + |b|$ and $(a^2 + b^2)^{1/2}$.

By suitable choices of \mathbf{u}_1 and \mathbf{u}_2 , we can read off various inequalities for $J(a, b)$. Firstly, $J(a, b) \leq J(a, 0) + J(0, b) = a + b$, as seen in (6). Secondly:

Second proof of (8). Since $(a + b, a + b) = (a, b) + (b, a)$, we have

$$(a + b) \frac{\pi}{2} = J(a + b, a + b) \leq J(a, b) + J(b, a) = 2J(a, b). \quad \square$$

Thirdly, we can derive an upper bound that simultaneously improves on those in (5) and (6), and is exact at both ends, at the cost of being unsymmetrical:

PROPOSITION 5. *For $0 \leq b \leq a$, we have*

$$J(a, b) \leq a + \left(\frac{\pi}{2} - 1\right) b. \quad (11)$$

Equality holds when $b = 0$ and when $b = a$.

Proof. Since $(a, b) = (a - b, 0) + (b, b)$,

$$J(a, b) \leq J(a - b, 0) + J(b, b) = (a - b) + \frac{\pi}{2} b. \quad \square$$

This bound is more natural than one might at first think. For fixed a , it represents the linear function of b that agrees with $J(a, b)$ at $b = 0$ and $b = a$. In fact, (11) reflects the fact that $J(a, b)$ is a *convex* function of b . Recall that a function is “convex” if it lies below the straight-line chords between pairs of points of its graph. This means that if $b_1 < b_2$ and $b_\lambda = (1 - \lambda)b_1 + \lambda b_2$, where $0 < \lambda < 1$, then

$$f(b_\lambda) \leq (1 - \lambda)f(b_1) + \lambda f(b_2).$$

By Proposition 4 and (E3), this holds with $f(b) = J(a, b)$, since $(a, b_\lambda) = (1 - \lambda)(a, b_1) + \lambda(a, b_2)$.

Comparison of the bounds, and a question about the derivative

Our two lower bounds for $J(a, b)$ are $m_1 = \frac{\pi}{4}(a + b)$ (exact when $b = a$) and $m_2 = (a^2 + b^2)^{1/2}$ (exact when $b = 0$). To explore how they compare, first fix $a = 1$. The two bounds coincide when $1 + b^2 = \frac{\pi^2}{16}(1 + b)^2$. The solutions of this quadratic are b_1 and $1/b_1$, where $b_1 \approx 0.34823$. The better (i.e. larger) lower bound is m_2 when $0 < b < b_1$ and when

$b > 1/b_1$. For general a , apply this to b/a to conclude that m_2 is the larger one when either b/a or a/b is less than b_1 .

Meanwhile, for $b \leq a$, our upper bounds are $M_1 = a + (\frac{\pi}{2} - 1)b$ (exact at both ends) and $M_2 = \frac{\pi}{2\sqrt{2}}(a^2 + b^2)^{1/2}$ (exact when $b = a$). When $a = 1$, the bounds coincide when $[1 + (\frac{\pi}{2} - 1)b]^2 = \frac{\pi^2}{8}(1 + b)^2$. One solution is $b = 1$, and the other is $b_2 \approx 0.25741$. The better (i.e. smaller) upper bound is M_1 when $\frac{b}{a} < b_2$.

Fixing $a = 1$, write $J_1(b)$ for $J(1, b)$. Do we now have enough information to give an accurate sketch of $J_1(b)$ as a function of b on the interval $[0, 1]$? The reader may care to draw a sketch of our four bounds on this interval, and it will be seen that between them they confine $J_1(b)$ to a fairly narrow box. But one ingredient for the sketch is still missing: it would be desirable to know the derivative, at least at the end-points. At $b = 1$, we do know: $J_1(b)$ is sandwiched between $m_1 = \frac{\pi}{4}(1 + b)$ and $M_2 = \frac{\pi}{2\sqrt{2}}(1 + b^2)^{1/2}$, and both have gradient $\pi/4$ there, hence $J_1'(1) = \pi/4$. Of course, this means that both m_1 and M_2 give good approximations to $J_1(b)$ for b close to 1. Also, m_1 is the tangent to the graph at $b = 1$, and the fact that the function is above the tangent again reflects convexity.

However, at $b = 0$, $J_1(b)$ is sandwiched between $(1 + b^2)^{1/2}$ (with gradient 0) and $1 + (\frac{\pi}{2} - 1)b$ (with gradient $\frac{\pi}{2} - 1$), which tells us nothing about $J_1'(0)$. We will now settle this problem. Contrary to what one might expect, we will do so by deploying some corresponding estimations for $I(a, b)$.

Inequalities for $I(a, b)$

A pair of bounds for $I(a, b)$ follows at once from identity (3): since $(ab)^{1/2} \leq M(a, b) \leq \frac{1}{2}(a + b)$, we have

$$\frac{\pi}{a + b} \leq I(a, b) \leq \frac{\pi}{2(ab)^{1/2}}. \quad (12)$$

The right-hand inequality can also be proved directly by the Cauchy-Schwarz inequality for integrals, most easily with $I(a, b)$ expressed in the equivalent form

$$I(a, b) = \int_0^\infty \frac{1}{(x^2 + a^2)^{1/2}(x^2 + b^2)^{1/2}} dx,$$

which equates to (2) by the substitution $x = b \tan \theta$.

The left-hand inequality in (12) does no justice to the fact that $I(a, b)$ actually tends to infinity as $b \rightarrow 0^+$. In fact, there is a theorem describing $I(1, b)$ very well for small b : a simple version of it states

$$\log \frac{4}{b} < I(1, b) < \log \frac{4}{b} + \frac{1}{2}b. \quad (13)$$

The proof, which is not very hard, can be seen in [Jam].

The integral $L(a, b)$, and a closer estimation for small b

Integrals of the following type form a very neat link between $J(a, b)$ and $I(a, b)$ (here we are following the account in [Lo]). For $a > 0$ and $b \geq 0$, define

$$L(a, b) = \int_0^{\pi/2} \frac{\cos^2 \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2}} d\theta. \quad (14)$$

The first thing to note is that $L(b, a) \neq L(a, b)$. In fact, the substitution $\theta = \frac{\pi}{2} - \phi$ gives

$$L(b, a) = \int_0^{\pi/2} \frac{\sin^2 \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{1/2}} d\theta. \quad (15)$$

From (14) and (15), we deduce at once (for $a, b > 0$) the following attractive pair of identities:

$$L(a, b) + L(b, a) = I(a, b), \quad (16)$$

$$a^2 L(a, b) + b^2 L(b, a) = J(a, b). \quad (17)$$

By eliminating $L(b, a)$ in (16) and (17), we deduce

$$(a^2 - b^2)L(a, b) = J(a, b) - b^2 I(a, b). \quad (18)$$

It is clear that $L(a, b)$ decreases with a and b , also $L(a, 0) = 1/a$ and $L(a, a) = \pi/(4a)$, so for $0 \leq b \leq a$, $L(a, b)$ lies between these two values. Note that $L(0, b)$ is not defined.

By differentiation under the integral sign, we have $\frac{\partial}{\partial b} J(a, b) = bL(b, a)$. To complete the symmetry, we mention that also $\frac{\partial}{\partial b} I(a, b) = -\frac{1}{b}L(a, b)$ (but this only becomes apparent after applying the substitution $x = b \tan \theta$ to both integrals).

So $J'_1(b) = bL(b, 1)$, and in particular, $J'_1(1) = L(1, 1) = \pi/4$, as we saw earlier. This expression for $J'_1(b)$ is undefined at $b = 0$, but we don't need it: we can easily determine $J'_1(0)$ using (18) and the elementary inequality (12).

PROPOSITION 6. *We have $J(1, b) \leq 1 + \frac{\pi}{2}b^{3/2}$ for $0 \leq b \leq 1$, hence $J'_1(0) = 0$.*

Proof. By (18),

$$J(1, b) = b^2 I(1, b) + (1 - b^2)L(1, b).$$

Now $L(1, b) \leq 1$ for $0 \leq b \leq 1$, so, with (12), we have $J(1, b) \leq 1 + b^2 I(1, b) \leq 1 + \frac{\pi}{2}b^{3/2}$. Hence

$$0 \leq \frac{J_1(b) - 1}{b} \leq \frac{\pi}{2}b^{1/2}.$$

This tends to 0 when $b \rightarrow 0^+$, which shows that $J'_1(0) = 0$. □

This might seem good enough. However, given (13), it will now cost us very little further work to derive corresponding bounds for $J(1, b)$, giving a highly accurate estimation for small b . We will express $J_1(b)$ as the integral of its derivative $bL(b, 1)$, so we want bounds for $L(b, 1)$.

LEMMA. For $0 < b < 1$,

$$\log \frac{1}{b} + c_1 \leq L(b, 1) \leq \log \frac{1}{b} + c_2, \quad (19)$$

where $c_1 = \log 4 - 1$ and $c_2 = \log 4 + \frac{1}{2} - \pi/4$.

Proof. By (16), we have $L(b, 1) = I(1, b) - L(1, b)$. Now $\frac{\pi}{4} \leq L(1, b) \leq 1$, and by (13),

$$\log \frac{1}{b} + \log 4 \leq I(1, b) \leq \log \frac{1}{b} + \log 4 + \frac{1}{2}$$

for $0 < b < 1$. Together, these inequalities give (19). \square

PROPOSITION 7. For $0 < b < 1$,

$$J(1, b) = 1 + \frac{1}{2}b^2 \log \frac{1}{b} + r(b), \quad (20)$$

where $c_3b^2 \leq r(b) \leq c_4b^2$, with $c_3 = \log 2 - \frac{1}{4}$ and $c_4 = \frac{1}{2} + \log 2 - \frac{\pi}{8}$ (so, in particular, $0 < r(b) < b^2$).

Proof. We have

$$J_1(b) - 1 = \int_0^b J'_1(t) dt = \int_0^b tL(t, 1) dt,$$

and by (19), $-t \log t + c_1t \leq tL(t, 1) \leq -t \log t + c_2t$. Now

$$\int_0^b (-t \log t) dt = \left[-\frac{1}{2}t^2 \log t\right]_0^b + \int_0^b \frac{1}{2}t dt = \frac{1}{2}b^2 \log \frac{1}{b} + \frac{1}{4}b^2.$$

The stated bounds are found by adding, respectively, $\int_0^b c_1t dt = \frac{1}{2}c_1b^2$ and $\frac{1}{2}c_2b^2$. \square

So we now have a pair of bounds that differ by less than b^2 . The previous lower bound $m_2 = (1 + b^2)^{1/2}$ has the correct gradient 0 at $b = 0$, but the lower bound in (20) is obviously stronger for small b . In fact, it is greater than $1 + \frac{1}{2}b^2$ (hence greater than m_2) for all $b \leq \frac{3}{4}$.

Since $J(a, 1) = aJ(1, \frac{1}{a})$, we can derive the following estimation, effective for large a :

$$J(a, 1) = a + \frac{\log a}{2a} + r_1(a),$$

where $c_3/a \leq r_1(a) \leq c_4/a$.

Finally, a fact that may seem rather surprising: J_1 does not have a second derivative at 0, because by the Lemma, $\frac{1}{b}J'_1(b) = L(b, 1) \rightarrow \infty$ as $b \rightarrow 0^+$.

References

- [Jam] G.J.O. Jameson, An approximation to the arithmetic-geometric mean, *Math. Gaz.* **98** (2014).
- [Lo] Nick Lord, Recent calculations of π ; the Gauss-Salamin algorithm, *Math. Gaz.* **76** (1992), 231–242.
- [LS] Nick Lord and H.-J. Seiffert, Solution to Problem 76D, *Math. Gaz.* **77** (1993), 126.