

INFINITE BAR-JOINT FRAMEWORKS, CRYSTALS AND OPERATOR THEORY

J.C. OWEN AND S.C. POWER

ABSTRACT. A theory of flexibility and rigidity is developed for general infinite bar-joint frameworks (G, p) . Determinations of nondeformability through vanishing flexibility are obtained as well as sufficient conditions for deformability. Forms of infinitesimal flexibility are defined in terms of the operator theory of the associated infinite rigidity matrix $R(G, p)$. The matricial symbol function of an abstract crystal framework is introduced, being the matrix-valued function on the d -torus representing $R(G, p)$ as a Hilbert space operator. The symbol function is related to infinitesimal flexibility, deformability and isostaticity. Various generic abstract crystal frameworks which are in Maxwellian equilibrium, such as certain 4-regular planar frameworks, are proven to be square-summably infinitesimally rigid as well as smoothly deformable in infinitely many ways. The symbol function of a three-dimensional crystal framework determines the infinitesimal wave flexes in models for the low energy vibrational modes (RUMs) in material crystals. For crystal frameworks with inversion symmetry it is shown that the RUMs appear in surfaces, generalising a result of F. Wegner [32] for tetrahedral crystals.

CONTENTS

1. Introduction	2
2. Infinite Bar-joint Frameworks	7
2.1. Continuous flexibility and rigidity	7
2.2. Linkages	10
2.3. Relative rigidity and the extension of flexes.	13
2.4. Forms of flexibility	17
2.5. Sufficient conditions for flexibility	18
2.6. Flex extensions and generic rigidity	19
3. Rigidity Operators and Infinitesimal Rigidity	20
3.1. Infinitesimal rigidity and the rigidity matrix.	20
4. Crystal Frameworks and Flexibility	25
4.1. Periodic and crystal frameworks.	26
4.2. Deformability and flow flexibility.	28
4.3. Crystal frameworks and periodic infinitesimal flexibility.	30

1991 *Mathematics Subject Classification.* 52C75, 46T20.

Key words and phrases. infinite bar-joint framework, vanishing flexibility, rigidity operator.

5. The Matricial Symbol Function of a Crystal Framework	32
5.1. Matrix function multiplication operators.	32
5.2. The rigidity matrix as a multiplication operator	33
5.3. From motifs to matrix functions.	35
5.4. Symmetry equations for infinite frameworks.	37
5.5. From matrix function to wave modes	38
5.6. Honeycomb frameworks and the kagome net.	41
5.7. Crystallography and Rigid Unit Modes.	44
References	45

1. INTRODUCTION

Infinite bar-joint frameworks appear frequently as idealised models in the analysis of deformations and vibration modes of amorphous and crystalline materials. See [11], [18], [6], [14], [32] and [36] for example and the comments below. Despite these connections there has been no extended mathematical analysis of such models. Notions of rigidity, flexibility, deformability, constrainedness, independence and isostaticity, for example, are usually employed either in the sense of their usage for a finite approximating framework or in a manner drawn from experience and empirical fact in the light of the application at hand. It seems that a deeper understanding of the models is of considerable interest in its own right and that a mathematical development may prove useful in certain applications. In what follows we shall provide formal definitions of the terms above in quite a wide variety of forms and we examine some of their inter-relationships and manifestations.

Suppose that one starts with a flexible square bar-joint framework in two dimensions and that this is then extended periodically to create an infinite periodic bar-joint network. Is the resulting assemblage, with inextendible bars, continuously flexible in two dimensional space? A moment's reflection reveals a proliferation of flexibility, such as sheering motions with one half of the network fixed.

However such movement is dramatically infinite and a natural second question is whether for periodic frameworks such as these there are flexes for which the total joint movement is finite. The less obvious negative answer in these cases offers some satisfaction in that it is consistent with so-called Maxwell counting in the case where the average number of degrees of freedom of the joints matches the average number of constraints per joint. (See Theorem 5.2 and Corollary 5.3.)

On the other hand if, for the grid example, one rigidifies alternate squares by adding diagonal bars, as shown, then the resulting structure of corner linked rigid squares remains properly flexible, although now uniquely so. In fact the unique flex has an affinely contracting character (See Definition 4.3) with alternating rotation of the squares.

More generally the flexibility of polytope networks in two and three dimensions continues to be of interest in the modeling of crystals and amorphous materials, especially with regard to their low frequency vibrational modes. Such modes appear, for example, in higher order symmetry phases of tetrahedral crystals and are referred to as rigid unit modes (RUMs). Indeed in the paper of Giddy et al [11] the alternating flex of the squares framework above has been associated with vibrational modes in perovskite. See also Hammond et al [18], Wegner [32], as well as Goodwin et al [14] for a useful overview. At the same time, in the modeling of amorphous materials, such as glasses, there is interest in understanding the critical probabilities that guarantee flexibility and rigidity for classes of randomly constructed frameworks. See, for example, Chubynsky and Thorpe [6] for the recent determination of such probabilities in simulation experiments.




FIGURE 1. The grid framework in the plane, $\mathcal{G}_{\mathbb{Z}^2}$.




FIGURE 2. The corner-joined squares framework, \mathcal{G}_{sq} .

Formally, a *framework* in \mathbb{R}^d (or bar-joint framework, or distance-constraint framework) is a pair $\mathcal{G} = (G, p)$ where $G = (V, E)$ is a simple connected graph and $p = (p_1, p_2, \dots)$ is a framework vector made up of framework points p_i in \mathbb{R}^d associated with the vertices v_1, v_2, \dots of V . The *framework edges* are the (closed) line segments $[p_i, p_j]$ associated with the edges E of the graph $G = (V, E)$. As the ellipsis suggest, we allow G to be a countable graph. We shall also define a *crystal framework* \mathcal{C} as a framework with translational symmetry which is generated by a connected finite *motif* of edges and vertices. (See Definition 4.2.)

When G is finite and the framework points are generically located in \mathbb{R}^2 then a celebrated theorem of Laman [21], well-known in structural engineering and in the discrete mathematics of rigidity matroids [15], gives a simple combinatorial criterion for the minimal infinitesimal rigidity of the framework; the graph itself satisfies Maxwell's counting rule $2|V| - |E| = 3$, and subgraphs $G' = (V', E')$ must comply with $2|V'| - |E'| \geq 3$. This is a beautiful result since the rigidity here is the noncombinatorial requirement that the kernel of an associated rigidity matrix $R(G, p)$ has the smallest dimension (namely three) for some (and hence all) generic framework. On the other hand frameworks with global symmetries, or even with "symmetric elements" (such as parallel edges) are not generic, that is, algebraic dependencies do exist between the framework point coordinates. Such frameworks arise in classical crystallography on the one hand and in mathematical models in structural engineering and in materials science on the other. See, for example, Donev and Torquato [7], Hutchinson and Fleck [19], Guest and Hutchinson [16] and various papers in the conference proceedings [31].

The present paper develops two themes. The first concerns a mathematical theory of deformability and rigidity for general infinite frameworks, with frequent attention to the case of periodic frameworks. There is, unsurprisingly, a great diversity of infinite framework flexing phenomena and we introduce strict terminology and some methods from functional analysis to capture some of this. In the second theme we propose an operator theory perspective for the infinitesimal (first order) flexibility of infinite frameworks.

Particularly interesting classes of infinite frameworks, from the point of view of flexibility, are those in the plane whose graphs are 4-regular and those in three dimensions whose graphs are 6-regular. In this case the graphs are in Maxwell equilibrium, so to speak, and so in a generic framework realisation any flex must activate countably many vertices. This is also the case for various periodic realisations such as the kagome framework, \mathcal{G}_{kag} , formed by corner-joined triangles in regular hexagonal arrangement, and frameworks in three dimensions formed by pairwise corner-joined tetrahedra. Despite being internally rigid in this way (Definition 2.18 (vi)) these frameworks admit diverse deformations. For example we note that the kagome framework admits uncountably many distinct deformations and in Theorem 4.4 we note that \mathbb{Z}^d -periodic cell-generic grid frameworks in \mathbb{R}^d admit deformations associated with affine transformations.

A significant phenomenon in the infinite setting is the appearance of *vanishing flexibility*. This means, roughly speaking, that the framework is a union of finite flexible subframeworks but the extent of flexibility diminishes to zero as the size of these subframeworks increases, so that

the infinite assemblage is inflexible. Elementary examples were indicated in [25] but we give more subtle examples here which are due to flex amplification at second order distances through concatenation effects. In particular there are bounded infinitesimal flexes in periodic frameworks that admit no continuous extensions and which do not arise as the derivative of a smooth deformation. We also note that there are \mathbb{Z}^2 -periodic crystal frameworks which are somewhat paradoxical, being indeformable despite the flexibility of all supercell subframeworks. On the other hand, in the positive direction, in Theorem 2.20 we give a general result which identifies a uniform principle for the existence of a deformation. The proof uses the Ascoli-Arzelà theorem on the precompactness of equicontinuous families of local flexes. It remains an interesting open problem to determine necessary and sufficient conditions for the rigidity and bounded rigidity of periodic planar frameworks.

The operator theory perspective for frameworks was suggested in [25] as an approach to a wider understanding of infinitesimal flexibility and rigidity. In this consideration the rigidity matrix is infinite and determines operators between various normed sequence spaces associated with nodes and with edges. Also, in [27] we have given a direct proof of the Fowler-Guest formula [9] for symmetric finite frameworks which is based on the commutation properties of the rigidity matrix as a linear transformation and this adapts readily to the infinite case and the rigidity operators of crystal frameworks. Indeed, translational symmetry ensures that the rigidity matrix $R(G, p)$ intertwines the coordinate shift operations. We consider square summable flexes and stresses and for distance regular locally finite frameworks $R(G, p)$ is interpreted as a bounded operator between Hilbert spaces. Also, enlarging to complex Hilbert spaces the Fourier transform $\mathcal{F}R(G, p)\mathcal{F}^{-1}$ is identified as a multiplication operator

$$M_\Phi : L^2(\mathbb{T}^d) \otimes \mathbb{C}^n \rightarrow L^2(\mathbb{T}^d) \otimes \mathbb{C}^m$$

given by an $m \times n$ matrix-valued function $\Phi(z)$ on the d -torus. The function Φ for \mathcal{C} is referred to as the *matricial symbol function* associated with the particular generating motif. The terminology and notation is borrowed from standard usage for Toeplitz operators and multiplication operators (see [3] for example). Many aspects of infinitesimal flexibility and isostaticity are expressible and analysable in terms of the matricial symbol function and its associated operator theory. For example a straightforward consequence of the operator theoretic approach is the square summable isostaticity of various non-degenerate regular frameworks that satisfy Maxwell counting, such as grid frameworks and the kagome framework.

An explicit motif-to-matrix function algorithm is given for the progression

$$\mathcal{C} = (G, p) \rightarrow R(G, p) \rightarrow \Phi(z).$$

Furthermore the identification of infinitesimal periodic-modulo-phase flexes and their multiplicities is determined by the degeneracies of $\Phi(z)$ as z ranges on the d -torus. In particular, the function

$$\mu(z) := \dim \ker \Phi(z) : \mathbb{T}^d \rightarrow \mathbb{Z}.$$

gives a determination of the mode multiplicity of periodic-modulo-phase infinitesimal flexes.

In the discussions below we are mainly concerned with properties of mathematical bar-joint frameworks. (The framework bars are indestructibly inextensible, the joints are located deterministically, they maintain perfect, frictionless fit and may even coincide.) Nevertheless, analysis of matricial symbol functions and their degeneracies are particularly relevant to the description and analysis of Rigid Unit Modes in material crystals. We show that for crystal frameworks with inversion symmetry the set of RUMS is a union of surfaces. This generalises and provides an alternative perspective for a recent result of Wegner [32] for tetrahedral crystals.

Operator theory methods have proven beneficial in many areas of mathematics and applications. This is evidently the case for multilinear systems theory and in control theory for example. Infinite rigidity matrix analysis seems to possess some similitudes with these areas and it seems to us that here too the operator turn will be a useful one.

The development is as follows. Section 2 gives a self-contained account of continuous flexibility and rigidity, and vanishing flexibility and one-sided flexibility is proven for various periodic infinite frameworks. Forms of flexibility, such as bounded flexes, square summable flexes, summable flexes and vanishing flexes are defined and determined for some specific examples. Sufficient conditions are obtained for the existence of a smooth flex and a flex extension problem for generic finite frameworks is posed. A positive resolution of this problem would provide a natural extension of Laman's theorem to infinite frameworks. In Section 3 we consider infinitesimal theory for general infinite frameworks and determine a number of rigidity operators and their flex and stress spaces. (The topic is taken up in more detail for crystal frameworks in Section 5.) In Section 4 we consider (abstract) crystal frameworks in two or three dimensions. These are generated by a motif and a discrete translation group. Various forms of deformations are considered, such as strict periodic flexibility, flow-periodic flexibility and flexes with reduced periodicity and symmetry. Also we indicate the flat torus model for crystal frameworks and recent results of Ross and Whiteley in this direction, including a periodic analogue of Laman's theorem. In the final section we consider the matricial symbol function approach and various examples. In particular we determine the (unit cell) infinitesimal wave flex multiplicities for the kagome net framework by factoring the determinant of the matricial symbol function.

Acknowledgments. Some of the developments here have benefited from discussions and communications with Robert Connelly, Patrick Fowler, Simon Guest, Elissa Ross and Walter Whiteley during and following the Summer Research Workshop on "Volume Inequalities and Rigidity", organised by Károly Bezdek, Robert Connelly, Balázs Csikős and Tibor Jordán, at the Department of Geometry, Institute of Mathematics, Eotvos Lorand University in July 2009.

2. INFINITE BAR-JOINT FRAMEWORKS

In this section we give a self-contained rigorous development of infinite frameworks and examine the nonlinear aspects of their flexibility by continuous deformations and their associated rigidity. In the next section we consider infinitesimal flexibility and rigidity in a variety of forms.

2.1. Continuous flexibility and rigidity. We first define continuous flexes and continuous rigidity. The latter means, roughly speaking, that the framework admits no proper deformations that preserve the edge lengths. The definition below gives straightforward generalisations of terms used for finite frameworks. In that case we note that a continuous flex is often referred to as a finite flex while in engineering models it is referred to as a finite mechanism.

Unless we specify otherwise we shall assume that the frameworks under consideration are *proper* in that the framework points do not lie on a hyperplane in the ambient space \mathbb{R}^d and that the framework edges $[p_i, p_j]$ have nonzero lengths $|p_i - p_j|$.

Definition 2.1. Let (G, p) be an infinite framework in \mathbb{R}^2 , with connected abstract graph $G = (V, E)$, $V = \{v_1, v_2, \dots\}$ and $p = (p_1, p_2, \dots)$.

(a) A base-fixed continuous flex, or, simply, a flex of (G, p) , is a function

$p(t) = (p_1(t), p_2(t), \dots)$ from $[0, 1]$ to $\prod_V \mathbb{R}^2$ with the following properties;

- (i) $p(0) = p$,
- (ii) each coordinate function $p_i : [0, 1] \rightarrow \mathbb{R}^2$ is continuous,
- (iii) for some *base edge* (v_a, v_b) with $|p_a - p_b| \neq 0$, $p_a(t) = p_a(0)$ and $p_b(t) = p_b(0)$ for all t ,
- (iv) each edge distance is conserved; $|p_i(t) - p_j(t)| = |p_i(0) - p_j(0)|$ for all edges (v_i, v_j) , and all t , and
- (v) $p(t) \neq p$ for some $t \in (0, 1]$.

(b) The framework (G, p) is *flexible*, or more precisely, *continuously flexible*, if it possesses a base-fixed continuous flex.

(c) The framework (G, p) is *rigid*, or continuously rigid, if it is not flexible.

Similarly one defines base-fixed continuous flexes and continuous rigidity for proper frameworks in \mathbb{R}^d by replacing a base edge by an appropriate set of framework points with maximal affine span.

The simplest kind of continuously rigid framework in the plane is one which is a union of continuously rigid finite frameworks. In particular the following theorem follows simply from the theorem of Laman indicated in the introduction.

Theorem 2.2. *Let G be a connected graph which is the union of a sequence of finite Laman graphs. Then every generic realisation (G, p) in the plane is continuously rigid.*

By *generic*, or, more precisely, *algebraically generic*, we mean, as is usual, that the coordinates of any finite set of framework points is algebraically independent over the rational numbers. Unlike the case of finite frameworks it is possible to construct two generic frameworks with the same abstract graph one of which is flexible and one of which is rigid. Accordingly it seems appropriate to formulate the following definition to extend the usual usage.

Definition 2.3. An infinite simple connected graph G is said to be rigid, or generically rigid, for two dimensions, if every generic framework (G, p) in the plane is rigid.

Note that if G is rigid and if H is a containing graph for G with the same vertex set then every generic framework (H, p) in the plane is rigid.

It seems likely that the converse to the theorem above holds. That is, if H does not contain a *sequentially Laman* graph (in the sense below) with the same vertex set, then H is not generically rigid. We comment more on this later in Section 2.6.

Rigidity and flexibility are properties of the entire framework and it is such entire features and their inter-relationships that are of primary interest in what follows. One would like to understand the relationship with small scale or local structure, such as local counting conditions and local connectivity. Additionally, as above, one would like to relate entire properties to sequential features that pertain to an exhausting chain of finite subframeworks and for this the following definition is helpful.

Definition 2.4. If P is a property for a class of finite, simple, connected graphs (resp. frameworks) then a graph G (resp. framework $\mathcal{G} = (G, p)$) is *sequentially P* or σ - P if G is the union of graphs in some increasing sequence of vertex induced finite subgraphs $G_1 \subseteq G_2 \subseteq \dots$, and each graph G_k (resp. framework (G_k, p)) has property P .

For example, we may refer to an infinite graph as being σ -Laman, or σ -Laman-1 and an infinite framework as being σ -rigid. To say that an

infinite framework is σ -flexible, or sequentially flexible, is rather vacuous since it usually prevails for trivial reasons. (One can construct countably infinite edge complete frameworks that may fail to be so but all our examples are sequentially flexible.) The more interesting property is the failure of sequentially rigidity as in the following definition.

Definition 2.5. A framework \mathcal{G} is said to have *vanishing flexibility* if it is continuously rigid but not σ -rigid.

An important topic in finite rigidity frameworks is that of *global rigidity*, also called unique rigidity. This holds, in the two dimensional setting, if there is, up to congruency, only one embedding of the framework in \mathbb{R}^2 . This implies, for example, that these frameworks admit no foldings, and indeed have unique diagrams up to rotations and reflection. One can extend the term to infinite frameworks but we do not consider this issue here at all. One might be tempted to say that a rigid infinite framework, especially one with vanishing flexibility, is globally rigid, but we refrain from doing so because of conflict with this usage.

The term "global" for global rigidity is natural since rigidity for finite frameworks is equivalent to the "local" property that "nearby equivalent frameworks are congruent". That is, if there exists $\epsilon > 0$ such that if (G, p') is a finite framework with $|p_i - p'_i| < \epsilon$, for all i , and if (G, p') is *equivalent* to (G, p) , in the sense that corresponding edges have the same length, then (G, p') and (G, p) are congruent. See, for example, Gluck [12] and Asimow and Roth [1].

An infinite simple graph G is *locally finite* if for every vertex v there are finitely many incident edges. Amongst such graphs are those for which there is an upper bound to the degree of the vertices, as in the case of the graphs of crystal frameworks. Within this class a graph G is said to be *r -regular* if every vertex has degree r . We remark that the theory of tilings provides a wealth of examples of planar frameworks which are 4-regular.

Remark 2.6. In what follows we consider only locally finite frameworks. Without this assumption it is possible to construct quite wildly flexing planar linkages. In fact, given a continuous function $f : [0, 1] \rightarrow \mathbb{R}^2$ one can construct an infinite linkage, in the sense of the definition below, and a base-fixed flex $p(t)$ with a motion $p_v(t)$ for a particular vertex v that is equal to $f(t)$. This includes the possibility of space filling curves. This is a consequence of a continuous analogue of a well known theorem of Kempe which asserts that any finite algebraic curve in the plane can be simulated by a finite linkage. For more details see Owen and Power [26].

2.2. Linkages. The removal of a framework edge from a rigid framework may result in flexibility which is, roughly speaking, of a one-dimensional nature. We reserve the term *linkage* for such a mathematical object, which we formally specify in the next definition. We remark that finite frameworks are also referred to as linkages, particularly when they are flexible, perhaps with several degrees of freedom, but this should not cause confusion.

A *two-sided continuous flex* $p(t)$ of (G, p) is defined as above but for the replacement of $[0, 1]$ by $[-1, 1]$. The following formal definition of an infinite linkage reflects the fact that the initial motion of a base-fixed linkage is uniquely determined by the angle change at any flexible joint.

Definition 2.7. A linkage in \mathbb{R}^2 is a finite or infinite connected framework $\mathcal{G} = (G, p)$ in \mathbb{R}^2 for which there exists a continuous two-sided base-fixed flex $p(t)$ with framework edges $[p_i, p_j], [p_j, p_k]$ such that the cosine angle function

$$g(t) = \langle p_i(t) - p_j(t), p_k(t) - p_j(t) \rangle$$

is strictly increasing on $[-1, 1]$, and such that $p(t)$ is the unique two-sided flex $q(t)$ of \mathcal{G} with $q_l(t) = p_l(t)$, for $l = i, j, k$.

Many interesting finite linkages were considered in the nineteenth century in connection with mechanical linkages. See, for example, Kempe [20]. Note however that the definition is liberal in allowing coincident joints and self-intersecting flexes. Also the definition refers to local deformation behaviour and this does not rule out the possibility of bifurcations occurring in a parameter extension of the given flex.

It is a simple matter to construct diverse infinite linkages by tower constructions or progressive assembly. (See, for example, the Cantor tree frameworks of [25].) However, some such constructions lead to frameworks with vanishing flexibility and so are not linkages in this case. An elementary illustration is given in Figure 4 wherein a two-way infinite rectangular strip linkage is augmented by adding flex-restricting cross braces in an alternating fashion. If the brace lengths tend to the diagonal length from above then the infinite framework is rigid. Evidently in this case the triangle inequality is playing a role in isolating one real solution to the *solution set* $V(G, p)$ defined below. One can also construct examples in which this isolation is less evident, with all joint angles bounded away from zero and $\pi/2$ for example.

A more interesting and subtle form of vanishing flexibility is due to progressive flex amplification rather than local flex restrictions. Roughly speaking, if a small flex is initiated at a particular joint and the flex propagates in some amplifying manner, then the triangle inequality at some far remove may prohibit any further increase. If the framework is infinite then no local joint flex may be possible at all. The strip

framework of concatenated levers in Figure 5 gives an example where the amplification is evident, while the rigidity of the strip framework in Figure 6 and the trapezium strip in Figure 7 is less evident. The lever framework has a natural infinitesimal flex, in the sense of Section 3, which is unbounded. The corresponding flexes of Figures 6 and 7 however, are bounded with amplification unfolding as a second order effect. This is proven in the next subsection.




FIGURE 3. An infinite rectangle strip linkage.




FIGURE 4. A restricted rectangle strip.




FIGURE 5. Rigid but not σ -rigid.




FIGURE 6. Rigid but not σ -rigid.

It is a straightforward matter to incorporate the vanishing flexibility of the strips above as subframeworks of a \mathbb{Z}^2 -periodic framework. This process is indicated in Figures 9, 10 and 11 below, where the infinite frameworks are determined as the periodic extensions of the given unit



FIGURE 7. Rigid but not σ -rigid.

FIGURE 8. A periodic half-strip which is only right-flexible.

cell. Figure 9 shows a linkage formed as a "fence lattice" composed of infinite horizontal and vertical σ -rigid bands. Figure 10 shows an analogue where the infinite bands have been replaced by rigid strip frameworks. Figure 11 is an elaboration of this in which cross braces have been introduced to remove the flexibility. (Only one edge is needed for this whereas the example given is periodic.) The additional degree 2 vertex in the cell ensures that the framework is not σ -rigid, while the infinite bands remain vanishingly rigid.

Let us note that for the framework in Figure 11, with its curious mixture of rigidity and flexibility, one can add any finite number of additional degree 2 vertices without changing the rigidity of the framework. In particular we have a construction that proves the following proposition.




FIGURE 9. Unit cell for a "fence lattice" linkage.

Proposition 2.8. *Let $c > 2$. Then there is a \mathbb{Z}^2 -periodic framework in \mathbb{R}^2 which is rigid, which is not σ -rigid and for which the average vertex degree is less than c .*





FIGURE 10. Unit cell for a modified fence lattice linkage.

FIGURE 11. Unit cell for a rigid periodic framework which is not σ -rigid.

One can readily extend this fanciful idea in various ways to obtain such structures in higher dimensions. For example, start with a one-dimensionally periodic σ -rigid girder in 3D and augment it with trapezium "tents" of alternating height to create vanishingly rigid girders. Also periodically interpolate any number of degree two vertices into the tent top edges without removing the vanishing flexibility. Join infinitely many such component girders periodically at appropriate tent-top edges to create a fence framework and add linear jointed cross braces to create, finally, a 2D periodic grid which is continuously rigid in 3D, which is not σ -rigid and which has average coordination number arbitrarily close to two.

2.3. Relative rigidity and the extension of flexes. For a finite or infinite framework $\mathcal{G} = (G, p)$ in \mathbb{R}^2 define the function

$$f_G : \prod_V \mathbb{R}^2 \rightarrow \prod_E \mathbb{R}, \quad f_G(q) = (|q_i - q_j|^2)_{e=(v_i, v_j)}.$$

This is the usual *edge function* of the framework and depends only on the abstract graph G .

Definition 2.9. The solution set, or configuration space, of a framework $\mathcal{G} = (G, p)$, denoted $V(G, p)$, is the set $f_G^{-1}(f_G(p))$. This is the set of all framework vectors q for G that satisfy the distance constraints

equations

$$|q_i - q_j|^2 = |p_i - p_j|^2, \text{ for all edges } e = (v_i, v_j).$$

In general the solution set of an infinite framework need not be a real algebraic variety even when it is "finitely parametrised". In less wild situations it can be useful to relate $V(G, p)$ to the algebraic variety $V(H, \pi_H(p))$ associated with a finite subgraph H of G , or with an elementary subgraph such as a tree, or even a set of vertices.

Definition 2.10. An infinite bar-joint framework (G, p) in \mathbb{R}^d is rigid over a subframework $(H, \pi_H(p))$ if every continuous flex of (G, p) which is constant valued on $(H, \pi_H(p))$ is constant. Similarly, if H is a subgraph of a countable connected simple graph G then G is *rigid over* H , or generically rigid over H if, for every generic frameworks (G, p) , every continuous flex of (G, p) which is constant-valued on $(H, \pi_H(p))$ is constant.

We may also form the following associated notions.

Definition 2.11. An infinite framework (G, p) in \mathbb{R}^d is finitely determined if it is rigid over (H, p) for some finite subgraph H and is finitely flexible if it is flexible and finitely determined.

Finite flexibility in the sense above is a strong property in which paths from p in the solution set $V(G, p)$ are determined near p by the finite algebraic variety $V(H, p)$. Note that the term "infinitely flexible" is not appropriate to describe a flexible framework which is not finitely flexible since it is possible to construct linkages, in our formal sense, which are not finitely flexible. This is the case for the periodic framework in Figure 12 which is a linkage because of partial vanishing flexibility.




FIGURE 12. A linkage which is not finitely determined.

An important class of frameworks which appear in mathematical models are those that are distance regular.

Definition 2.12. A framework $\mathcal{G} = (G, p)$ in \mathbb{R}^d is distance-regular if there exist $0 < m < M$ such that for all edges (i, j) ,

$$m < |p_i - p_j| \leq M.$$

For such a framework (G, p) it is natural to consider the nearby frameworks with the same graphs but with slightly perturbed framework points (and therefore edge lengths). If a property holds for all such frameworks, for some perturbation distance ϵ then we call such a property a *stable property* for the framework.

Formally, an ϵ -perturbation of a distance regular framework $\mathcal{G} = (G, p)$ is a framework $\mathcal{G}' = (G, p')$ for which $|p_i - p'_i| < \epsilon$, for all corresponding framework points. Recall that a finite framework in \mathbb{R}^d is said to be ϵ -rigid if it is congruent to every *equivalent* ϵ -perturbation. Let us say that a general framework is *perturbationally rigid* if it is ϵ -rigid for some ϵ . It is a well-known fact that perturbational rigidity and rigidity are equivalent in the case of algebraically generic finite frameworks [1], [12]. However, it is straightforward to see that this equivalence thoroughly fails for general infinite frameworks (see [25]).

Definition 2.13. Let \mathcal{G} be a distance-regular framework. Then \mathcal{G} is stably rigid (resp stably flexible) if it is rigid (resp. flexible) and for sufficiently small $\epsilon > 0$ every ϵ -perturbation of \mathcal{G} is rigid (resp. flexible).

Likewise, if P is any particular property of a distance-regular infinite framework then we may say that \mathcal{G} is *stably P* if, for some $\epsilon > 0$, the property P holds for all ϵ -perturbations.

Proposition 2.14. *The periodic trapezium strip frameworks, with alternating unequal heights $a > b > 0$, are rigid. In particular the rectangle strip linkage (of Figure 7) is not stably flexible.*




FIGURE 13

Proof. Figure 13 shows the displacement of a double trapezium to the right. Let the three vertical bar lengths be a, b and a units with $a > b > 0$. The displaced position has angles A, B, C at the base line and angles D, E, F, G occurring relative to the trajectory tangents of the displaced vertices. For a subsequent incremental change δA , with resulting incremental changes $\delta B, \delta C, \delta D, \delta E, \delta F, \delta G$ we can see from simple geometry that to first order

$$a(\delta A \cos D) = b(\delta B \cos E).$$

Suppose now that A is regarded as a specialisation of the input angle α with resulting output angle $\beta = \beta(\alpha)$, so that at $\alpha = A$ we have $\beta(\alpha) = B$. Then

$$\frac{d\beta}{d\alpha}\Big|_{\alpha=A} = \lim_{\delta A \rightarrow 0} \frac{\delta B}{\delta A} = \frac{a \cos D}{b \cos E}.$$

Similarly, with angle C regarded as the output angle $\gamma(B)$ for the angle transmission function $\gamma = \gamma(\beta)$, we have

$$\frac{d\gamma}{d\beta}\Big|_{\beta=B} = \frac{b \cos F}{a \cos G}$$

and so

$$\frac{d\gamma}{d\alpha}\Big|_{\alpha=A} = \frac{d\gamma}{d\beta}\Big|_{\beta=B} \frac{d\beta}{d\alpha}\Big|_{\alpha=A} = \frac{\cos D \cos F}{\cos E \cos G}.$$

Note that since $B > A$ we have also $D > E$, and since $B > C$ we have $F > G$, from which it follows that both ratios above are less than one. Thus certainly $0 < \gamma'(\alpha) < 1$ for $0 < \alpha < \alpha_1$ where α_1 is the first positive angle for which $\gamma'(\alpha_1) = 0$.

It follows, from the mean value theorem, that the double trapezium angle transmission function is an increasing differentiable function with

$$\gamma(0) = 0, \quad 0 < \gamma(\alpha) < \alpha, \quad \text{for } 0 < \alpha < \alpha_1.$$

It follows immediately that the right-semi-infinite trapezium strip is right flexible.

We let $\lambda = \gamma(\alpha_1)$, which we refer to as the locking angle. Note the second trapezium of the double admits no increase of this angle. In view of the above we have $\lambda < \alpha_1$.

Suppose now that \mathcal{G} is the two-way-infinite trapezium strip, with $a \neq b$. Let $p(t)$ be a flex and suppose that for a fixed framework edge with length a the angle A is greater than zero for some time $t_1 > 0$ and that t_1 is the first such time. Then certainly $0 < A < \lambda$. Note that $A_{-n} = \gamma^{-n}(A)$, $n = 1, 2, \dots$ are the angles of the edges of length a , counted off to the left. In view of the function dominance $0 < \gamma(\alpha) < \alpha$ it follows that $A_{-n} > \alpha_1$ for some n , which is a contradiction. \square

The argument above also shows that the semi-infinite trapezium strip framework of Figure 7 has a continuous flex but no two-sided flex.

Remark 2.15. It seems to be of interest to analyse strip frameworks in further detail. For example, a trapezium strip framework is not stably rigid, despite the apparent "robustness" of the argument above. To see this use surgery in the following way. Remove one cross bar, then push the rightmost semifinite strip to the right, by an angle perturbation $A = \epsilon > 0$. Now insert a replacement bar of the required length. One can flex the resulting structure towards the left to restore the position of the right hand strip. Indeed, this is all the flexibility the framework has. The possible flex of an ϵ perturbation, such as the one described,

seems to be of order ϵ and so there does seem to be "approximate rigidity".

Remark 2.16. Consider a periodic trapezium grid framework \mathcal{G}_{trap} obtained by perturbing $\mathcal{G}_{\mathbb{Z}^2}$ by adding a fixed small positive value to the y coordinate of the framework points p_{ij} for the odd values of i and j . It can be shown that this framework is rigid over any linear subframework. This contrasts with the grid framework itself which is freely flexible over its x and y axes in the following specific local sense: all sufficiently small flexes of the subframework extends to a flex of the whole framework. On the other hand note that Theorem 4.4 shows that \mathcal{G}_{trap} is deformable.

2.4. Forms of flexibility. It seems to be a fundamental and interesting issue to determine the ways in which infinite bar-joint frameworks are rigid or continuously flexible. In this section we give some further definitions, we give sufficient conditions for the existence of a proper flex and we contemplate a plausible infinite framework version of Laman's theorem.

Flexes are often infinitely differentiable or smooth in the sense of the following formal definition. This is the case for example, for the "alternation" flexes of \mathcal{G}_{sq} and \mathcal{G}_{kag} .

Definition 2.17. A continuous base-fixed two-sided flex $p(t) : t \in [-1, 1]$ of a framework (G, p) in \mathbb{R}^d is a *smooth flex* if each coordinate function $p_i(t)$ is infinitely differentiable.

The smoothness of a flex is a local requirement whereas the following terms impose various increasing forms of global constraint. In particular rotational flexes of infinite frameworks with unbounded diameter are not bounded flexes, while a translational flex of an infinite framework is a bounded flex but is not a vanishing flex. Adopting a term that has been used in applications [14] we refer to flexes which are not bounded as *colossal flexes*.

Definition 2.18. A continuous flex $p(t) = (p_k(t))_{k=1}^{\infty}$, ($t \in [0, 1]$) of an infinite framework (G, p) in \mathbb{R}^d is said to be

- (i) a bounded flex if for some $M > 0$ and every k and t ,

$$|p_k(t) - p_k(0)| \leq M,$$

- (ii) a colossal flex if it is not bounded,

(iii) a vanishing flex if $p(t)$ is a bounded flex and if the maximal displacement

$$\|p_k - p_k(0)\|_{\infty} = \sup_{t \in [0, 1]} |p_k(t) - p_k(0)|$$

tends to zero as $k \rightarrow \infty$,

(iv) a square-summable flex if

$$\sum_{k=1}^{\infty} \|p_k - p_k(0)\|_{\infty}^2 < \infty,$$

(v) a summable flex if

$$\sum_{k=1}^{\infty} \|p_k - p_k(0)\|_{\infty} < \infty,$$

(vi) an internal flex if for all but finitely many k the function $p_k(t)$ is constant.

Also we say that (G, p) has a deformation (resp. bounded or vanishing deformation) if it has a base-fixed flex $p(t)$ (which is bounded or vanishing).

Definition 2.19. A connected infinite locally finite proper framework in two or three dimensions is boundedly rigid (resp. summably rigid, square-summably rigid, smoothly rigid, internally rigid) if there is no deformation, that is, no base-fixed proper continuous flex, which is bounded (resp. summable, square-summable, smooth, internal).

2.5. Sufficient conditions for flexibility. There is a sense in which vanishing flexibility is almost the only obstacle to the existence of a flex of a framework all of whose finite subframeworks are flexible. More precisely, in the hypotheses of the next theorem we assume that there are two distinguished framework vertices, p_1, p_2 , such that any finite subframework (H, p) containing p_1, p_2 has a flex which properly separates this pair in the sense of condition (ii) below. The additional requirement needed is that there is a family of flexes of the finite subframeworks whose restrictions to any given subframework (H, p) are uniformly smooth in the sense of condition (i). Note that the constant here depends only on (H, p) and indeed the resulting flex may of necessity be a colossal flex.

Theorem 2.20. *Let (G, p) be an infinite locally finite framework in \mathbb{R}^d with a connected graph, let*

$$(G_1, p) \subseteq (G_2, p) \subseteq \dots,$$

be subframeworks, determined by finite subgraphs $G_r = (V_r, E_r)$ with union equal to G and let v_1, v_2 be vertices in G_1 . Suppose moreover that for each $r = 1, 2, \dots$, there is a base-fixed smooth flex $p^r(t) = (p_k^r(t))_{k=1}^{|V(G_r)|}$ of $\mathcal{G}_r = (G_r, p)$ such that

(i) for each finite framework \mathcal{G}_l the set \mathcal{F}_l of restriction flexes

$$\{p^r(t)|_{\mathcal{G}_l} : r \geq l\}$$

have uniformly bounded derivatives, that is, there are constants $M_l, l = 1, 2, \dots$, such that

$$\left| \frac{d}{dt} p_k^r(t) \right| \leq M_l \quad \text{for } r \geq l, v_k \in V_l,$$

(ii) the framework points p_1, p_2 are uniformly separated by each flex $p^r(t)$ in the sense that

$$|p_1^r(1) - p_2^r(1)| - |p_1^r(0) - p_2^r(0)| \geq c$$

for some positive constant c .

Then (G, p) has a deformation.

Note that it is essential that the separated vertices of condition (ii) are the same for each subgraph. To see this note that the two-way infinite trapezium strip framework considered in Figure 7 has smooth deformations on each of its finite strip subframeworks, each of which "separates" some two vertices (at the end of the strip) by a fixed positive distance. Nevertheless the infinite strip fails to have a deformation.

Proof. For $l = 1, 2, \dots$, let X_l be the space of continuous functions from $[0, 1]$ to $\mathbb{R}^{d|V_l|}$ and note that the family \mathcal{F}_l , by the hypotheses, is an equicontinuous family in X_l . Moreover with respect to the supremum norm \mathcal{F}_l is a bounded set. By the Ascoli-Arzelà theorem (see [30] for example) \mathcal{F}_l is precompact and in particular for \mathcal{F}_1 there is a subsequence r_1, r_2, \dots such that the restrictions $p^{r_k}|_{\mathcal{G}_1}$ converge uniformly in their (finitely many) coordinates to a flex q^1 of \mathcal{G}_1 . Relabel the sequence (p^{r_k}) as $(p^{(k,1)})$. The restrictions of these flexes to \mathcal{G}_2 similarly have a convergent subsequence, say $(p^{(k,2)})$, and so on. From this construction select the diagonal subsequence $(p^{(k,k)})$. This converges coordinatewise uniformly to a coordinatewise continuous function

$$q : [0, 1] \rightarrow \mathbb{R}^d \times \mathbb{R}^d \times \dots$$

Since the restriction of q to every finite subframework is a flex, q satisfies the requirements of a flex of (G, p) , except possibly the properness requirement (v) of Definition 2.1. In view of (ii) however, $q(0) \neq q(1)$ and so q is a continuous flex of (G, p) . \square

Remark 2.21. Computer simulations provide evidence for the fact that small random perturbations of $\mathcal{G}_{\mathbb{Z}^2}$ yield frameworks that are flexible. That is it seems that $\mathcal{G}_{\mathbb{Z}^2}$ is stably flexible. It would be interesting if the theorem above could assist in a proof of this.

2.6. Flex extensions and generic rigidity. Let us recall a version of Laman's theorem.

Theorem 2.22. *Let (G, p) be an algebraically generic finite framework. Then (G, p) is infinitesimally rigid if and only if the graph G has a vertex induced subgraph H , with $V(H) = V(G)$, which is maximally*

independent in the sense that $2|V(H)| = |E(H)| + 3$ and $2|V(H')| \geq |E(H')| + 3$ for every subgraph H' of H .

For convenience we refer to a maximally independent finite graph as a Laman graph. We remark that any Laman graph can be obtained from a triangle graph by a sequence of moves known as Henneberg moves. The first of these adds a new vertex with two connecting edges while the second breaks an edge into two at a new vertex which is then connected by a new edge to another point of the graph.

Now let G be an infinite graph which contains a subgraph H on all the vertices of G and suppose that H is σ -Laman. In view of Laman's theorem every algebraically generic realisation of H (and hence G) in the plane is σ -rigid and so continuously rigid. Is the converse true? That is, if every generic realisation of an infinite graph G is rigid does G necessarily contain a σ -Laman subgraph with the same vertex set.

To see this one needs to show is that if G is σ -(Laman-1) and not σ -Laman, then there exists a vertex generic realisation (G, p) in the plane which has a continuous flex. That is, we want to build up a flex of the infinite structure by adding new vertices and edges, in the least handicapping way, to allow all, or most of the flex of an initial finite subgraph to be extended.

Alternatively, and more explicitly, suppose that one starts with a generic connected Laman-1 framework (G_1, p) with n vertices and an infinite sequence of Henneberg move "instructions". These instructions yield a unique infinite graph. Is it possible to choose associated framework points p_n, p_{n+1}, \dots so judiciously that some (perhaps small) flex of (G_1, p) extends fully to each successive finite extension framework (and hence the infinite framework)?

3. RIGIDITY OPERATORS AND INFINITESIMAL RIGIDITY

In previous sections we have considered some variety in the *nature* of continuous flexes $p(t)$ and how they might distinguished. A companion consideration is the analysis of various spaces of infinitesimal flexes. This gives insight into continuous flexes since the derivative $p'(0)$ of a differentiable flex $p(t)$ is an infinitesimal flex.

Here we give an operator theory perspective for an infinitesimal theory of infinite frameworks in which the rigidity matrix $R(G, p)$ is viewed as a linear transformation or linear operator between various spaces. The domain space contains a space of infinitesimal flexes, which lie in the kernel of the rigidity operator, while the range space contains a space of self-stress vectors namely those in the kernel of the transpose of $R(G, p)$.

3.1. Infinitesimal rigidity and the rigidity matrix. Recall that for a finite framework (G, p) in \mathbb{R}^d with $n = |V|$ an *infinitesimal flex* is a vector $u = (u_1, \dots, u_n)$ in the vector space $\mathcal{H}_v = \mathbb{R}^d \oplus \dots \oplus \mathbb{R}^d$

such that the orthogonality relation $\langle p_i - p_j, u_i - u_j \rangle = 0$ holds for each edge (v_i, v_j) . This condition ensures that if each p_i is perturbed to $p_i(t) = p_i + tu_i$, with t small, then the edge length perturbations are of second order only as t tends to zero. That is, for all edges,

$$|p_i(t) - p_j(t)| - |p_i - p_j| = O(t^2).$$

If $q(t) : [-1, 1] \rightarrow \mathcal{H}_v$ is a two-sided smooth flex of the finite framework (G, p) then $q'(0)$ is an infinitesimal flex and for a generic finite framework every infinitesimal flex arises in this way. See Asimow and Roth [1] for example.

Associate with an infinite framework (G, p) the product vector space

$$\mathcal{H}_v = \prod_V \mathbb{R}^d = \mathbb{R}^d \oplus \mathbb{R}^d \oplus \dots$$

consisting of all sequences $u = (u_1, u_2, \dots)$. Conceptually such a vector corresponds to a specification of instantaneous velocities, or to a perturbation sequence, applied to the framework joints. Define an *infinitesimal flex* of (G, p) to be a vector u in \mathcal{H}_v for which, as above, $\langle p_i - p_j, u_i - u_j \rangle = 0$ holds for each edge (v_i, v_j) , and let \mathcal{H}_{fl} denote the linear space of all these vectors. In the planar case \mathcal{H}_{fl} contains the three-dimensional linear subspace (assuming G has at least one edge) of the infinitesimal flexes that arise from the isometries of \mathbb{R}^2 . Note that a nonzero rotation infinitesimal flex u is an unbounded sequence if and only if (G, p) is an unbounded framework. We denote the space of rigid body motion infinitesimal flexes as \mathcal{H}_{rig} .

Definition 3.1. An infinite framework (G, p) is infinitesimally rigid if every infinitesimal flex is a rigid body motion infinitesimal flex.

If G is infinite then \mathcal{H}_v contains properly the direct sum space $\mathcal{H}_v^{00} = \sum_V \mathbb{R}^2$ consisting of vectors whose coordinates are finitely supported, in the sense of being finitely nonzero. The following definition is convenient and evocative.

Definition 3.2. An infinite framework (G, p) is internally infinitesimally rigid if every finitely supported infinitesimal flex is the zero flex.

We now give the usual direct definition of the rigidity matrix $R(G, p)$ of a framework (G, p) , allowing G to be infinite. This matrix could also be introduced via the Jacobian of the equation system that defines $V(G, p)$ since $2R(G, p)$ is the Jacobian evaluated at p .

Write $p_i = (x_i, y_i)$, $u_i = (u_i^x, u_i^y)$, $i = 1, 2, \dots$, and denote the coordinate difference $x_i - x_j$ by x_{ij} . The rigidity matrix is an infinite matrix $R(G, p)$ with rows indexed by edges e_1, e_2, \dots and columns labeled by vertices but with multiplicity two, namely $v_1^x, v_1^y, v_2^x, v_2^y, \dots$. Note that any matrix of this shape, with finitely many nonzero entries in each row, provides a linear transformation from \mathcal{H}_v to $\mathcal{H}_e = \prod_E \mathbb{R}$.




FIGURE 14. Unit cell for an internally infinitesimally flexible periodic framework.

Definition 3.3. The rigidity matrix of the infinite framework (G, p) , with $p = (p_i) = (x_i, y_i)$, is the matrix $R(G, p)$ with entries $x_{ij}, x_{ji}, y_{ij}, y_{ji}$ occurring in the row with label $e = (v_i, v_j)$ with the respective column labels $v_i^x, v_j^x, v_i^y, v_j^y$, and with zero entries elsewhere.

In particular a vector u in \mathcal{H}_v is an infinitesimal flex if and only if $R(G, p)u = 0$.

Definition 3.4. (i) A stress (or, more properly, a self-stress) of a finite or infinite framework (G, p) is a vector $w = (w_e)$ in $\mathcal{H}_e = \prod_E \mathbb{R}$ such that $R(G, p)^t w = 0$.

(ii) A finite or infinite framework (G, p) is isostatic, or absolutely isostatic, if it is infinitesimally rigid and has no nonzero self-stresses.

Since it is understood here that G is a locally finite graph a rigidity matrix has finitely many entries in each column and so its transpose corresponds to a linear transformation from \mathcal{H}_e to \mathcal{H}_v .

In the finite case a self-stress represents a finite linear dependence between the rows of the rigidity matrix, which one might abbreviate, with language abuse, by saying that the corresponding edges of the framework are linearly dependent. A self-stress vector $w = (w_e)_{e \in E}$ can be simply related to a vector $b = (b_e)$ conceived of as a sequence $b = (b_e)$ of bar tension forces with a resolution, or balance, at each node. Indeed, for such a force vector b the vector w for which $w_e = |p_i - p_j|^{-1} b_e$ ($e = (v_i, v_j)$) is a stress vector. Thus there is a simple linear relationship between the space of internal stresses and the space of resolving bar tensions. We shall not consider here the more general stress vectors, important in engineering applications, that arise from an external loading vectors.

Let \mathcal{H}_e^{00} be the space of finitely supported vectors in \mathcal{H}_e . We say that an infinite framework (G, p) is *finitely isostatic* if it is internally infinitesimally rigid and if the finite support stress space $\mathcal{H}_{str}^{00} := \mathcal{H}_{str} \cap \mathcal{H}_e^{00}$ is equal to $\{0\}$. It is straightforward to see that the grid frameworks $\mathcal{G}_{\mathbb{Z}^d}$, in their ambient spaces, are finitely isostatic, as is the kagome framework.

Between the extremes of infinitesimal rigidity and internal rigidity there are other natural forms of rigidity such as those given in the following definition. Write ℓ^∞ , ℓ^2 and c_0 to indicate the usual Banach sequence spaces for countable coordinates, and write $\mathcal{H}_e^\infty, \mathcal{H}_v^\infty, \dots, \mathcal{H}_v^0$ for the corresponding subspaces of \mathcal{H}_e and \mathcal{H}_v .

Definition 3.5. An infinite framework (G, p) is

(i) square-summably infinitesimally rigid (or infinitesimally ℓ^2 -rigid) if

$$\mathcal{H}_v^2 \cap \ker R(G, p) = \{0\},$$

(ii) boundedly infinitesimally rigid (or infinitesimally ℓ^∞ -rigid) if

$$\mathcal{H}_v^\infty \cap \ker R(G, p) = \mathcal{H}_v^\infty \cap \mathcal{H}_{rig},$$

(iii) vanishingly infinitesimally rigid (or infinitesimally c_0 -rigid) if

$$\mathcal{H}_v^0 \cap \ker R(G, p) = \{0\},$$

(iv) square-summably isostatic (or ℓ^2 -isostatic) if it is infinitesimally ℓ^2 -rigid and

$$\mathcal{H}_{str} \cap \mathcal{H}_e^2 = \{0\},$$

(v) boundedly isostatic if it is boundedly infinitesimally rigid and $\mathcal{H}_{str} \cap \mathcal{H}_e^\infty = \{0\}$,

(vi) vanishingly isostatic if it is vanishingly infinitesimally rigid and $\mathcal{H}_{str} \cap \mathcal{H}_e^0 = \{0\}$.

There is companion terminology for flexes and stresses. Thus we refer to vectors in $\mathcal{H}_v^2 \cap \ker R(G, p)$ as square summable infinitesimal flexes and so on.

Example 3.6. Let us use the shorthand (\mathbb{N}, p) to denote a semi-infinite framework in \mathbb{R}^2 whose abstract graph is a tree with a single branch, with edges $(v_1, v_2), (v_2, v_3), \dots$ and where $p = (p_i), p_i = (x_i, y_i), i = 1, 2, \dots$. Then, writing x_{ij} and y_{ij} for the differences $x_i - x_j$ and $y_i - y_j$, as before, the rigidity matrix with respect to the natural ordered bases takes the form

$$R(\mathbb{N}, p) = \begin{bmatrix} x_{12} & y_{12} & x_{21} & y_{21} & 0 & \dots & & \\ 0 & 0 & x_{23} & y_{23} & x_{32} & y_{32} & 0 & \dots \\ 0 & 0 & 0 & 0 & * & * & * & \dots \\ \vdots & & & & & & & \end{bmatrix}.$$

With respect to the coordinate decomposition $\mathcal{H}_v = \mathcal{H}_x \oplus \mathcal{H}_y$ we have

$$R(\mathbb{N}, p) = \begin{bmatrix} R_x & R_y \end{bmatrix} = \begin{bmatrix} D_x & D_y \end{bmatrix} \begin{bmatrix} T \\ T \end{bmatrix}$$

where $R_x = D_x T$, $R_y = D_y T$, where D_x and D_y are the diagonal matrices

$$D_x = \begin{bmatrix} x_{12} & 0 & 0 & \dots \\ 0 & x_{23} & 0 & \dots \\ 0 & 0 & x_{34} & 0 \\ \vdots & & & \ddots \end{bmatrix} \quad D_y = \begin{bmatrix} y_{12} & 0 & 0 & \dots \\ 0 & y_{23} & 0 & \dots \\ 0 & 0 & y_{34} & 0 \\ \vdots & & & \ddots \end{bmatrix}$$

and

$$T = \begin{bmatrix} 1 & -1 & 0 & \dots \\ 0 & 1 & -1 & \dots \\ 0 & 0 & 1 & -1 \\ \vdots & & & \ddots \end{bmatrix}.$$

If we now identify the domain and range spaces in the natural way for these coordinates then we have $T = I - U^t$ where U^t is the transpose of the forward unilateral shift operator on the linear space of real sequences.

The analogous framework (\mathbb{Z}, p) has a similar matrix structure in all respects except that in place of the Toeplitz matrix T one has the corresponding two-way infinite Laurent matrix $I - W^{-1}$ where W is the forward bilateral shift. In both cases, the two-dimensional subspace spanned by the translation flexes is evident, being spanned by the constant vectors in \mathcal{H}_x and \mathcal{H}_y . Evidently there are infinitely many finitely supported flexes and in fact it is possible to identify $\ker R(G, p)$ as a direct product space.

One can use operator formalism to examine the space of stresses. In the case of the simple framework (\mathbb{Z}, p) note that $W^t = W^{-1}$ and

$$\ker(I - W) = \ker(I - W^{-1}) = \mathbb{R}e$$

where e is the vector with every entry equal to 1. Since

$$R(\mathbb{Z}, p)^t = \begin{bmatrix} I - W & \\ & I - W \end{bmatrix} \begin{bmatrix} D_x \\ D_y \end{bmatrix}$$

it follows that a vector w is a stress vector if and only if $D_x w \in \mathbb{R}e$ and $D_y w \in \mathbb{R}e$. Thus for some constants α, β we have $x_{i,i+1} w_i = \alpha$, $y_{i,i+1} w_i = \beta$, and so for all i

$$\frac{y_i - y_{i+1}}{x_i - x_{i+1}} = \frac{\beta}{\alpha}.$$

This colinearity condition shows that the space of stresses is trivial unless the framework points p_i , $i \in \mathbb{Z}$ are colinear in which case \mathcal{H}_{str} is one dimensional. This includes the colinear cases in which p is a bounded sequence and the framework lies in a finite line segment in \mathbb{R}^2 .

Example 3.7. With similar notational economy write (\mathbb{Z}^r, p) , (resp. (\mathbb{N}^r, p)) for frameworks associated with the *grid graph* with vertex set

labeled by r -tuples of integers (resp. positive integers) $n = (n_1, \dots, n_r)$ where the edges correspond to vertex pairs $(n, n \pm e_j)$, where e_1, \dots, e_r are the usual basis elements. The ambient space for the framework is either understood or revealed by the entries of the vector p .

Again one can use operator formalism to analyse the space of stresses as a vector subspace of \mathcal{H} . In the special case of the regular grid framework $\mathcal{G}_{\mathbb{Z}^2}$ in \mathbb{R}^2 one can see that the vector subspace \mathcal{H}_{str} is a direct product vector space (like \mathcal{H} itself) whose product basis is indexed by (two-way infinite) linear subframeworks parallel to the coordinate axes. This is also true for a general *orthogonal grid framework* such as the bounded grid framework determined by the framework points $(\pm(1 - (1/2)^i), \pm(1 - (1/2)^j))$.

Note that we have defined an infinitesimal flex in a local way, being the verbatim counterpart of the usage for finite frameworks. In particular the notion takes no account of the possibility of (second order) amplification or vanishing flexibility.

Our examples above indicate the importance of shift operators and in the next section we see that the bilateral shift operators, in their Fourier transform realisation as multiplication operators, play a central role in the discussion of periodic frameworks.

Let us note here that approximate infinitesimal flexes are natural for infinite frameworks and the operator theoretic perspective.

Definition 3.8. An approximate square-summable flex of an infinite framework (G, p) is a sequence of finitely supported unit vectors u_1, u_2, \dots in \mathcal{H}_v^2 (or \mathcal{K}_v^2) such that $\|R(G, p)u_n\|_2 \rightarrow 0$ as $n \rightarrow \infty$.

Let (G, p) be a distance regular framework. Then it is straightforward to show that the rigidity matrix determines a bounded Hilbert space operator R . It is the metrical and geometric properties of the action of R and its transpose that have relevance to rigidity theory rather than the spectral theory of R . However we do have the almost vacuous statement that the existence of approximate (square-summable) flexes corresponds to the point 0 belonging to the approximate point spectrum of R . For if 0 lies in the approximate point spectrum then (by definition) Rv_n is a null sequence for some sequence of unit vectors v_n , and approximation of these unit vectors by vectors with finite support yields, after normalisations, an approximate square-summable flex sequence (u_n) .

It is implicit in the matricial function association below that the rigid unit modes of translationally periodic frameworks are tied to the existence of approximate flexes.

4. CRYSTAL FRAMEWORKS AND FLEXIBILITY

In previous sections we have constructed frameworks to illustrate various definitions and properties. It is perhaps of wider interest to

understand, on the other hand, how extant infinite frameworks, such as those suggested by crystals or repetitive structures, may be flexible. Accordingly we now define crystal frameworks and investigate various forms of flexibility and rigidity.

4.1. Periodic and crystal frameworks. We have already observed some properties of the basic examples of the *grid framework* $\mathcal{G}_{\mathbb{Z}^2}$, the *squares framework* \mathcal{G}_{sq} and the kagome framework, \mathcal{G}_{kag} . In \mathbb{R}^3 we also have analogues, such as the cube framework \mathcal{G}_{cube} , the octagon framework \mathcal{G}_{oct} and the kagome net framework \mathcal{G}_{knet} , which consist, respectively, of vertex-joined cubes, octahedra and tetrahedra, with no shared edges or faces, each in a natural periodic arrangement. The frameworks \mathcal{G}_{kag} , \mathcal{G}_{cube} , \mathcal{G}_{oct} and \mathcal{G}_{knet} are polytope body-pin frameworks but we consider the polytope rigid units as bar-joint subframeworks formed by adding some, or perhaps all, internal edges. As such these frameworks are examples of crystal frameworks in the sense of the formal definition below. We first comment on a wider notion of periodicity.

Definition 4.1. An affinely periodic framework in \mathbb{R}^d is a framework $\mathcal{G} = (G, p)$ for which there exists a non-trivial discrete group of affine transformations $T_g, g \in \mathcal{D}$, where each T_g acts on framework points and framework edges.

For example, the two-way infinite dyadic cobweb framework of Figure 15 is affinely periodic for the dilation doubling map and the four-fold dihedral group D_4 .




FIGURE 15. An affinely periodic cobweb framework.

For another example we may take the infinite \mathbb{Z} -periodic framework in three dimensions for which Figure 16 forms a perspective view down a central axis, with framework vertices $(\pm 1, \pm 1, m), m \in \mathbb{Z}$. Here the affine group is an isometry group isomorphic, as a group, to $\mathbb{Z} \times C_2 \times D_4$.

To illustrate the following definition observe in Figure 16 a template of six edges and three vertices which generates the kagome framework by the translations associated with the parallelogram unit cell. Borrowing crystallographic terminology we refer to such a template as a *motif* for the framework and the chosen translation group. The following formal definition gives a convenient way of specifying abstract crystal frameworks.




FIGURE 16. A motif and unit cell for the kagome framework.

Definition 4.2. A crystal framework $\mathcal{C} = (G, p)$ in \mathbb{R}^d is a connected bar-joint framework for which there is a discrete group of translation isometries $\mathcal{T} = \{T_g : g \in \mathcal{D}\}$, a finite connected set F_e of framework edges and a finite set F_v of framework vertices (being a subset of the vertex set of F_e) such that

(i) the unions

$$\cup_{g \in \mathcal{D}} T_g(F_e), \quad \cup_{g \in \mathcal{D}} T_g(F_v)$$

coincide with the sets of framework edges and vertices, and

(ii) these unions are unions of disjoint sets.

We denote a crystal framework \mathcal{C} by the triple (F_v, F_e, \mathcal{T}) or by the triple (F_v, F_e, \mathbb{Z}^d) in the case of integer translations. An associated *unit cell* for \mathcal{C} may be defined as a set which contains F_v and no other framework points and for which the translates under the translation isometries are disjoint and partition the ambient space. For example in the case of $\mathcal{G}_{\mathbb{Z}^2}$ we may take the semiopen set $[0, 1)^2$ or the set $[0, 1) \times [1/2, 5/2)$ as unit cells. Such parallelepiped unit cells are useful for us for torus models for crystal frameworks. Voronoi cells (Brillouin zones) also play a unit cell role in applications but we shall not need such geometric detail here.

In many applied settings the appropriate framework models have "short" edges, spanning no more than two adjacent unit cells. Here we allow general edges which may span a chain of adjacent cells.

Recall from elementary crystallography that, modulo orthogonal transformations, there are 14 different forms (or symmetry types) in which a countable set of isolated points can be arranged with translational symmetry throughout three-dimensional space. These arrangements

are called the Bravais lattices and the translation group \mathcal{T} above corresponds to such a lattice. Thus each point of the framework lies in the Bravais lattice generated by the orbit of its unique corresponding motif vertex under the translational group.

4.2. Deformability and flow flexibility. Recall that a general (countable, locally finite, connected) framework \mathcal{G} is *rigid* if there is no base-fixed continuous flex and is *boundedly rigid*, or boundedly non-deformable, if there is no bounded base-fixed continuous flex $p(t)$. Recall, from Section 2, that bounded flexes are those for which there is an absolute constant M such that for every vertex v the time separation $|p_v(t) - p_v(0)|$ is bounded by M for all t and all v .

We first describe a context for the standard "alternation" flexes of \mathcal{G}_{sq} and \mathcal{G}_{kag} and certain periodic flexes of \mathcal{G}_{kag} with reduced symmetry. For these nonbounded flexes translational periodicity is maintained but relative to an affine flow of the ambient space. By an *affine flow* we mean simply a path $t \rightarrow A_t$ of affine transformations of \mathbb{R}^d which is pointwise continuous. The simplest such flow in two dimensions is a contracting flow such as $A_t(x, y) = ((1 - tc)x, (1 - tc)y)$, $0 < c < 1$. The alternating flexes of \mathcal{G}_{sq} and \mathcal{G}_{kag} are associated with such a flow.

Definition 4.3. Let $d = 2, 3$, let $t \rightarrow A_t$ be a flow of \mathbb{R}^d and let $\mathcal{C} = (G, p)$ be a crystal framework for the translation group $\mathcal{T} = \{T_g : g \in \mathcal{D}\}$. A flow-periodic flex of \mathcal{C} , relative to the flow and the translation group \mathcal{T} , is a continuous flex $p(t)$ such that for each t the framework $(G, p(t))$ is \mathcal{T}^t -periodic, where $\mathcal{T}^t = \{A_t T_g A_t^{-1} : g \in \mathcal{D}\}$.

Let us say simply that \mathcal{C} is *affinely periodically deformable* if there exists a non trivial flow-periodic flex.

A flow-periodic flex $p(t)$ for \mathcal{C} can be defined for a given flow if and only if for each t one can continuously solve the distance constraint equations for the vertex positions of the motif, with the periodicity constraint, in the A_t -deformed unit cell. Equivalently, the motif and unit cell define a finite framework on a torus with the noninterior edges of the motif providing reentrant ("locally geodesic") edges on the torus. For example consider the torus framework in Figure 17.




FIGURE 17. A torus framework.

A horizontal affine contraction $H_t : (x, y) \rightarrow ((1 - t)x, y)$ leads to a continuous flex of the torus framework and hence to a colossal flex of the associate crystal framework. The same is true for the vertical affine contraction V_t and for the skew affine transformation

$$S_t : (x, y) \rightarrow (x + (\sin t)y, y + (1 - \cos t)x).$$

Note that this particular transformation preserves both the cyclic width and cyclic height of the torus.

For an illustration of the method we note the following affine deformation result for what might be termed periodic cell-generic grid frameworks in \mathbb{R}^d . For more general results of this nature see also Borcea and Streinu [2].

Theorem 4.4. *Let $\mathcal{C} = (\mathbb{Z}^d, p)$ be a grid framework in \mathbb{R}^d which is (n_1, \dots, n_d) -periodic and which is a $1/3$ -perturbation of $\mathcal{G}_{\mathbb{Z}^d}$ in the sense that*

$$|p(k_1, \dots, k_d) - (k_1, \dots, k_d)| < 1/3$$

for $0 \leq k_i \leq n_i - 1, 1 \leq i \leq d$. Then \mathcal{C} is affinely periodically deformable.

Proof. We sketch the proof in case $d = 2$ with $(n_1, n_2) = (n, m)$. For convenience re-scale the framework so that the large cell $[0, n - 1] \times [0, m - 1]$ of \mathcal{C} becomes to the usual unit cell. Let (F_v, F_e, \mathbb{Z}^2) be the motif representation of (\mathbb{Z}^2, p) . Suppose first that $n = m = 2$ and the motif has four noninterior edges (corresponding to the four reentrant edges on the associated torus). Let F'_e be F_e with one vertical and one horizontal reentrant edge removed and let \mathcal{C}' be the associated framework. (See Figure 17.) For t taking positive values in some finite interval there are affine deformations of \mathcal{C}' associated with each of the flows H_t, V_t, S_t . Moreover there is an affine deformation associated with any composition $A_t = V_{\beta(t)}H_{\alpha(t)}S_t$, where the functions α and β are any continuous functions with $\alpha(0) = \beta(0) = 0$, where t takes values in some finite interval. Now note that we may chose $\alpha(t)$ so that the separation distance corresponding to the omitted horizontal edge is constant, and we may choose $\beta(t)$ similarly for the omitted vertical edge. Note also the essential fact that $V_{\beta(t)}$ does not change the cyclic width. In this way an affine deformation for \mathcal{C} is determined. (In the case of generic points, is determined uniquely on some finite interval).

The same principle operates for general (n, m) . The subframework \mathcal{C}' , with two deleted noninterior edges (one for each coordinate direction) once again has a flow periodic deformation associated with a composition $A_t = V_{\beta(t)}H_{\alpha(t)}S_t$, for t in a sufficiently small interval. This follows from a simple induction argument. Once again the functions

α and β are chosen to provide a deformation of \mathcal{C}' in which the separations for the omitted edges is constant. In this way a flow periodic deformation of \mathcal{C} itself is determined. \square

The deformations obtained in the last proposition are colossal deformations and this seems to be a necessary condition if some form of periodicity is to be maintained. It would be interesting to determine when such frameworks possess bounded "unstructured" deformations.

One may also consider affine deformations relative to a subgroup of \mathcal{D} associated with a supercell. By a *supercell* we mean a finite union (or possibly an infinite union) of adjacent cells which tile the ambient space by translations from the subgroup. In this case the deformations maintain only a longer period of translational symmetry and for an infinite linear supercell one may even forgo translational symmetry in one direction.

To illustrate this let us note a class of interesting deformations of the kagome framework (G, p) which have this form.

Start with an alternation flex $p(t)$ which leaves fixed a particular vertex p_* and leaves invariant a line of hexagon diameters. Note that this flex (which is not base-fixed) has bilateral symmetry with respect to the line (viewed as a mirror line) but that the inversion symmetry of (G, p) about the fixed vertex is broken for $t > 0$. Perform identical surgeries on the frameworks $(G, p(t))$ as follows:

- (i) cut the frameworks along the fixed line,
- (ii) effect a reflection of one of the resulting half-planes frameworks in the orthogonal line through p_* and rejoin the half-planes to create a new flex of $(G, p(0))$.

We might view this flex as one with a symmetry transition line. The hexagons in this line maintain inversion symmetry while elsewhere the hexagons maintain bilateral symmetry.

One may perform such surgery on several parallel surgery lines simultaneously. Performing surgery on countably many lines leads to the following theorem. This shows, roughly speaking, that the kagome framework sits in its configuration space as an infinitely singular point in the sense that it is the starting point for uncountably many distinct flexes. (Here we use the term distinct in the sense of Definition 2.7.)

Theorem 4.5. *There are uncountably many distinct flow periodic flexes of the kagome framework.*

4.3. Crystal frameworks and periodic infinitesimal flexibility.

Let $\mathcal{C} = (G, p) = (F_v, F_e, \mathbb{Z}^2)$ be a crystal framework in the plane for the integer translation isometry group. A natural form of infinitesimal flex for \mathcal{C} is that of a 1-cell periodic flex in the following sense.

Definition 4.6. A vector $u = (u_v)_{v \in V}$ in the real vector space \mathcal{H}_v , or in the complex vector space \mathcal{K}_v , is a 1-cell-periodic infinitesimal flex for

\mathcal{C} (or simply a periodic flex if there is no ambiguity) if $u \in \ker R(G, p)$ and $u_{\kappa+n} = u_\kappa$ for all $\kappa \in F_v$ and $n \in \mathbb{Z}^d$.

Such an infinitesimal flex u is a bounded sequence of vectors determined by periodic extension of what we may call the *motif flex vector* $u_{motif} = (u_\kappa)_{\kappa \in F_v}$. There is a one-to-one correspondence between these periodic flexes and the finite vectors that are in the kernel of the *motif rigidity matrix* $R_m(G, p)$ which we may define as the natural "periodic completion" of the $|F_e| \times d|F_v|$ submatrix of $R(G, p)$. This is the natural representing matrix for $R(G, p)$ viewed as a linear transformation between the finite dimensional subspaces of \mathbb{Z}^2 -periodic vectors. If there are no reflexive edges in the motif then

Similarly we define periodic infinitesimal stresses as those which correspond to periodic extensions of vectors in the cokernel of the motif rigidity matrix.

The crystal framework \mathcal{C} , with given motif and discrete translation group indexed by \mathcal{D} , is said to be *1-cell-periodically isostatic* (or simply periodically isostatic) if the only 1-cell periodic flexes are translation flexes and if there are no nontrivial 1-cell periodic stresses.

Once again it can be helpful to consider a flat torus model for such frameworks. Note for example that the generic periodic framework defined by Figure 17 is periodically rigid and indeed periodically isostatic.

The following interesting periodic variant of Laman's theorem has been obtained recently by E. Ross [28]. Let us say that the planar periodic framework \mathcal{C} is topologically proper if \mathcal{C} is connected and for every torus subframework motif F'_v, F'_e with $2|F'_v| - |F'_e| = 2$ there is an edge cycle from F'_e which properly wraps around the torus in the sense that the associated homotopy class is nonzero.

Theorem 4.7. *Let $\mathcal{C} = (F_v, F_e, \mathbb{Z}^2)$ be an infinite framework in the plane which is periodic for the integer translation group and is topologically proper. Then following are equivalent.*

(i) $2|F_v| - |F_e| = 2$ and for all edge induced submotifs F'_v, F'_e we have $2|F'_v| - |F'_e| \geq 2$.

(ii) \mathcal{C} is periodically isostatic in the sense that the periodic vectors in $\ker R(G, p)$ are spanned by the two translations (periodic rigidity) and the periodic vectors in $\text{coker } R(G, p)$ are zero (periodic stress free).

As we have seen, a periodically infinitesimally rigid framework in the sense above may nevertheless correspond to a flat torus framework which can flex infinitesimally (and even deform) relative to contraction or expansion of the unit cell. Allowing such vertical and horizontal freedoms adds two further degrees of freedom to the constraint equations. It would be of interest to obtain a similar characterisation in this case as well as for the triply flexible affine torus.

5. THE MATRICIAL SYMBOL FUNCTION OF A CRYSTAL FRAMEWORK

We now derive matrix function operators for general (abstract) crystal frameworks. Our approach is decidedly Hilbert space theoretic and allows for an extended conceptual framework for rigidity analysis. First we outline the standard identification of operators commuting with shift operators and multiplication operators in a Fourier transform space. We adopt complex scalars, replacing the \mathcal{H}_v spaces of Section 3 by their complex counterparts, $\mathcal{K}_v, \mathcal{K}_e, \mathcal{K}_v^\infty, \mathcal{K}_v^2$, and so forth.

5.1. Matrix function multiplication operators. Let ϕ be a continuous complex-valued function on the unit circle \mathbb{T} . It defines a multiplication operator T on the usual complex Hilbert space $L^2(\mathbb{T})$. Its representing matrix with respect to the standard orthonormal basis $\{z^n : n \in \mathbb{Z}\}$ is the $\mathbb{Z} \times \mathbb{Z}$ indexed matrix with entries

$$T_{i,j} = \langle \phi z^j, z^i \rangle = \langle \phi z^{j-i}, 1 \rangle = \hat{\phi}(i-j),$$

the $(i-j)^{th}$ Fourier coefficient of ϕ .

Similarly, let $\Phi(z)$ be a continuous matrix-valued function on the two or three dimensional torus \mathbb{T}^d taking values in the space of $n \times m$ complex matrices. One can specify such a function $\Phi(z) = \Phi(z_1, \dots, z_d)$ in terms of a matrix of scalar functions $\Phi(z) = [\phi_{k,l}(z)]_{k=1, l=1}^{n,m}$. Given in this way $\Phi(z)$ defines a multiplication operator from $L^2(\mathbb{T}) \otimes \mathbb{C}^m$ to $L^2(\mathbb{T}) \otimes \mathbb{C}^n$. Indeed, choose a basis $\{\xi_l\}$ for \mathbb{C}^m and an associated basis $\{\xi_l \otimes z^p\}$ for the domain space. Similarly let $\{\eta_k \otimes z^p\}$ be a basis for the codomain. Then the operator T of multiplication by $\Phi(z)$ may be defined by specifying, for each continuous function $f(z)$ on \mathbb{T}^d ,

$$T(\xi_l \otimes f(z)) = \sum_{k=1}^m \eta_k \otimes \phi_{k,l}(z) f(z),$$

and extending by linearity and continuity. Thus we can see that the representing matrix for T with respect to these bases is determined by the Fourier coefficients of the matrix entries for Φ :

$$\langle T(\xi_l \otimes z^p), \eta_k \otimes z^q \rangle_{\mathbb{C}^m \otimes L^2(\mathbb{T}^d)} = \langle \phi_{k,l}(z) z^{p-q}, 1 \rangle_{L^2(\mathbb{T}^d)} = \hat{\phi}_{k,l}(q-p).$$

Viewing an element of $L^2(\mathbb{T}^d) \otimes \mathbb{C}^m$ as a function $F(z)$ taking values in \mathbb{C}^m (strictly speaking, taking values almost everywhere), and similarly for vectors in the codomain, one also considers TF as the function $\Phi(z)F(z)$. This operator is usually denoted as M_Φ .

Suppose now, starting afresh, we have orthonormal bases

$$\{\xi_{l,p} : 1 \leq l \leq m, p \in \mathbb{Z}^d\} \quad \{\eta_{k,p} : 1 \leq k \leq n, p \in \mathbb{Z}^d\}$$

for Hilbert spaces such as \mathcal{K}_v^2 and \mathcal{K}_e^2 respectively. Suppose also that we are given an operator T from \mathcal{K}_v^2 to \mathcal{K}_e^2 by means of its representing matrix with respect to these bases and suppose moreover that this

matrix has the translational symmetry above in the sense that the matrix entries

$$\langle T\xi_{l,p}, \eta_{k,q} \rangle_{\mathcal{K}_e^2}$$

are independent of $p - q$. Let $\mathcal{F}_v : \mathbb{C}^n \otimes L^2(\mathbb{T}^d) \rightarrow \mathcal{K}_v^2$ be the Fourier transform and \mathcal{F}_e the Fourier transform for \mathcal{K}_e^2 . These are simply the unitary operators determined by the natural bijection of basis elements, namely $\xi_{l,p} \rightarrow \xi_l \otimes z^p$ and $\eta_{k,p} \rightarrow \eta_k \otimes z^p$, respectively. Thus, from the scalar matrix entries the functions $\phi_{k,l}$ are determined and hence the matrix function $\Phi(z)$. In this way one identifies $\mathcal{F}_e^{-1}T\mathcal{F}_v$ as the associated multiplication operator \mathcal{M}_Φ .

5.2. The rigidity matrix as a multiplication operator. Let us keep in view the special case of the \mathbb{Z}^2 -periodic grid framework determined by an algebraically generic quadrilateral in the unit cell.

Let p^1, \dots, p^4 be four framework vertices in the unit cell $[0, 1]^2$ constituting the motif set F_v of \mathbb{Z}^2 -periodic quadrilateral grid (\mathbb{Z}^2, p) . Let $p_{ij}^t = p^t + (i, j)$, for $(i, j) \in \mathbb{Z}^2$, be the general framework points and let e^1, \dots, e^8 denote the eight edges which form the motif set F_e given by

$$e^1 = [p^1, p^2], e^2 = [p^2, p^3], e^3 = [p^3, p^4], e^4 = [p^4, p^1],$$

$$e^5 = [p_{-1,0}^2, p^1], e^6 = [p_{0,-1}^3, p^2], e^7 = [p_{1,0}^4, p^3], e^8 = [p_{0,1}^1, p^4],$$

and let $e_{i,j}^t = e^t + (i, j)$ be a typical edge. Write the corresponding basis for the vector space \mathcal{K}_e^0 as

$$\{\eta_{t,i,j} : 1 \leq t \leq 8, (i, j) \in \mathbb{Z}^2\}.$$

Also, label the natural basis for \mathcal{K}_v^0 as the set of vectors $\{\xi_{s,i,j}^x, \xi_{s,i,j}^y\}$, where s ranges from 1 to 4 and (i, j) range through \mathbb{Z}^2 .

The rigidity matrix $R(G, p)$ satisfies the following symmetry equations,

$$W_{i,j}R(G, p)U_{i,j} = R(G, p)$$

with respect to the translation shift operators $U_{i,j}$ on \mathcal{K}_v^{00} and $W_{i,j}$ on \mathcal{K}_v^{00} . By the discussion above the rigidity matrix for this framework gives rise to an 8×8 matrix function $\Phi(z_1, z_2) = (\phi_{i,j}(z_1, z_2))$ on the torus \mathbb{T}^2 in \mathbb{C}^2 which is determined by the equations

$$\hat{\phi}_{k,\{x,s\}}(-i, j) = \langle (R(G, p)\xi_{s,i,j}^x, \eta_{k,0,0}) \rangle,$$

together with a companion set of equations for the y -labeled basis elements. Furthermore there is a simple algorithm for computing the symbol matrix function from the motif (F_v, F_e) and the motif rigidity matrix.

For a general crystal frameworks in the plane we have the following recipe for identifying $\Phi(z_1, z_2)$. There is an entirely similar identification of the matrix function of a crystal framework in higher dimensions.

Theorem 5.1. *Let $\Phi(z)$ be the matricial symbol function of the crystal framework $\mathcal{C} = (F_v, F_e, \mathbb{Z}^2)$. The entry of the rigidity matrix determined by the edge e_k of F_e and the column labeled $v_{s,i,j}^x$ (resp. $v_{s,i,j}^y$) provides the $(-i, -j)$ -th Fourier coefficient of $\phi_{k,\{x,s\}}$ (resp. $\phi_{k,\{y,s\}}$).*

Note that for the motif for the periodic quadrilateral grid each edge has vertices with different s index. This is rather typical and in such cases it follows that $\phi_{k,\{x,l\}}$ is either zero or has one nonzero Fourier coefficient. In the case of a "reflexive" edge in F_e , of the form $[p_{\kappa,0}, p_{\kappa,\delta}]$ there are entries in the columns for κ corresponding to $(1 - \bar{z}^\delta)v_e$ where v_e is the usual vector of coordinate differences appearing in the rigidity matrix for the edge e . That is, $v_e = p_{\kappa,0} - p_{\kappa,\delta}$.

Thus we may obtain the following identification where the direct construction of Φ from \mathcal{C} is given above.

Theorem 5.2. *Let $\mathcal{C} = (F_v, F_e, \mathbb{Z}^d)$ be a crystal framework with $m = |F_e|$, $n = |F_v|$ and with rigidity operator R from \mathcal{K}_v^2 to \mathcal{K}_e^2 .*

(i) *The motif (F_v, F_e) determines a matrix-valued trigonometric function $\Phi : \mathbb{T}^d \rightarrow M_{m,dn}(\mathbb{C})$ for which there is a unitary equivalence $\mathcal{F}_e^{-1}R\mathcal{F}_v = M_\Phi$ where M_Φ is the multiplication operator*

$$M_\Phi : \mathbb{C}^{nd} \otimes L^2(\mathbb{T}^d) \rightarrow \mathbb{C}^m \otimes L^2(\mathbb{T}^d).$$

(ii) *The framework \mathcal{C} is square-summably rigid (resp. square-summably stress-free) if the column rank (resp. row rank) of $\Phi(z)$ is maximal for almost every $z \in \mathbb{T}^d$.*

The following corollary shows that the existence of a square summable flex is a rather strong condition for a crystal framework and in fact "typical" frameworks in Maxwell counting equilibrium have no such flexes. What is of particular significance for such frameworks however is the presence of approximate flexes, in the sense noted in Section 3, as these are related to the local rank degeneracies of $\Phi(z)$ and to periodic flexes.

Corollary 5.3. *The following are equivalent for a crystal framework with Maxwell counting equilibrium.*

- (i) *\mathcal{C} has a nonzero internal (finitely supported) infinitesimal flex.*
- (ii) *\mathcal{C} has a nonzero summable infinitesimal flex.*
- (iii) *\mathcal{C} has a nonzero square-summable infinitesimal flex.*

Proof. Note that (i) implies (ii) and (ii) implies (iii) so it remains to show (iii) implies (i).

Suppose that the matrix $\Phi(z)$ has a square-summable vector $f(z)$ in its kernel. Then $\det(\Phi(z)) = 0$ for almost all points in the support of $f(z)$. Since $\det \Phi(z)$ is a multivariable trigonometric polynomial it necessarily vanished identically, that is the polynomial $\det \Phi(z)$ is the zero polynomial.

Recall that for any square matrix X over a ring we have $X\tilde{X} = \det(X)I_n$ where \tilde{X} is given by the usual formula $\tilde{X}_{i,j} = (-1)^{i+j}X_{i,j} \det(X^{ij})$ involving the cofactors (X^{ij}) . In particular if \tilde{X} is not zero and $\det(X) = 0$ then for a nonzero column vector $f(z)$ of \tilde{X} we have $Xf(z) = 0$. Applying this to $X = \Phi$ provides the desired polynomial (finitely supported) vector in this case.

If \tilde{X} happens to be the zero matrix we make use of the minimal polynomial lemma below. To apply the lemma let q be the minimum polynomial of X so that $q(X) = 0$. By the lemma we have $q(X) = Xq_1(X)$ and $q_1(X) \neq 0$ by minimality. Thus there is a nonzero vector $f(z)$ in the range of $q_1(X)$ and this provides the desired local flex. \square

Lemma 5.4. *Let $q(\lambda)$ in $R[\lambda]$ be the minimal polynomial of an n by n matrix X whose entries lie in an integral domain R and suppose that $\det X = 0$. Then $q(0) = 0$.*

Proof. Let $p(\lambda)$ be the characteristic polynomial $\det(\lambda I_n - X)$. Recall from the Cayley-Hamilton theorem that $p(X) = 0$. In the algebraic closure of the field of fractions of R the linear factors of $q(\lambda)$ and $p(\lambda)$ agree. See Lange [22] for example. Thus

$$q(\lambda) = \prod (\lambda - \alpha_i)^{s_i}, \quad p(\lambda) = \prod (\lambda - \alpha_i)^{r_i}$$

with $s_i \leq r_i$ for each i . In particular for suitably large n the polynomial $q(0)^n$ is divisible by $p(0)$. Since $\det X = 0$ it follows that $p(0) = 0$ and so $q(0)^n = 0$. Since $q(0)$ is in R it is equal to zero and the proof is complete. \square

5.3. From motifs to matrix functions. We now consider some examples. Adjusting notation, write z, w for the coordinates variables of \mathbb{T}^2 so that a trigonometric polynomial takes the form a finite sum

$$\phi(z, w) = \sum_{i,j} \hat{\phi}(i, j) z^i w^j.$$

Examining the recipe of Theorem 5.1 above in the case of the periodic quadrilateral grid we see that the internal edges e^1, \dots, e^4 of the motif provide four rows for $\Phi(z, w)$ whose entries are constant functions with constants corresponding to those of the rows for the rigidity matrix $R(G, p)$. The other edges provide rows according to the following simplified monomial rule. As we have intimated above, this rule applies generally when all the "external" edges of the motif have vertices with distinct s index:

the entries of the e^{th} row of $\Phi(z, w)$ that appear in the columns for the external (non-motif) vertex are the corresponding entries for the motif rigidity matrix multiplied by the monomial $\bar{z}^i \bar{w}^j$, where (i, j) is the shift index for the cell occupied by the external vertex.

Thus we obtain the matrix identification in the next proposition. We write $p_i = (x_i, y_i)$, for $i = 1, \dots, 4$. Note, for example, that the entry $x_{12} + 1 = x_1 - (x_2 - 1)$ denotes the constant function corresponding to the entry of $R(G, p)$ for row e^5 and column x_1 , while $-(x_{12} + 1)z$ is the entry corresponding to the x -coordinate of the external vertex of e^5 .

Proposition 5.5. *The matricial symbol function $\Phi(z, w)$ of the periodic quadrilateral grid framework (\mathbb{Z}^2, p) for the four vertex motif is the matrix function on \mathbb{T}^2 given by*

$$\begin{bmatrix} x_{12} & y_{12} & -x_{12} & -y_{12} & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{23} & y_{23} & -x_{23} & -y_{23} & 0 & 0 \\ 0 & 0 & 0 & 0 & x_{34} & y_{34} & -x_{34} & -y_{34} \\ -x_{41} & -y_{41} & 0 & 0 & 0 & 0 & x_{41} & y_{41} \\ x_{12} + 1 & y_{12} & -(x_{12} + 1)z & -y_{12}z & 0 & 0 & 0 & 0 \\ 0 & 0 & x_{23} & y_{23} + 1 & -x_{23}w & -(y_{23} + 1)w & 0 & 0 \\ 0 & 0 & 0 & 0 & x_{34} - 1 & y_{34} & (1 - x_{34})\bar{z} & -y_{34}\bar{z} \\ -x_{41}\bar{w} & (1 - y_{41})\bar{w} & 0 & 0 & 0 & 0 & x_{41} & y_{41} - 1 \end{bmatrix}$$

Furthermore, if the four points in the unit cell have algebraically independent coordinates then (\mathbb{Z}^2, p) is square-summably isostatic.

Proof. The first four rows are linearly independent since the points p_1, \dots, p_4 have algebraically independent coordinates. It can be shown from elementary linear algebra for almost every z, w the full matrix has rank 8. Thus Theorem 5.2 applies. \square


As a special case we obtain the matrix symbol function for $\mathcal{G}_{\mathbb{Z}^2}$ determined by the eight-edged motif, namely

$$\Phi(z, w) = 1/2 \begin{bmatrix} -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & -z & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & -w & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & \bar{z} & 0 \\ 0 & \bar{w} & 0 & 0 & 0 & 0 & 0 & -1 \end{bmatrix}$$

The determinant in this case is

$$-\frac{1}{256} zw(1 - \bar{z})^2(1 - \bar{w})^2.$$

On the other hand for this symmetric quadrilateral framework $\mathcal{G}_{\mathbb{Z}^2}$ the formalism above can be applied to a smaller unit cell with the simple two-edged motif of Figure 18.

FIGURE 18. A motif and unit cell for $\mathcal{G}_{\mathbb{Z}^2}$.

Note that the index s for the recipe takes the value 1 only and both motif edges have vertices with the same s index. It follows that the rigidity operator of $\mathcal{G}_{\mathbb{Z}^2}$ associated with the motif is unitarily equivalent to the multiplication operator on the Hilbert space $\mathbb{C}^2 \otimes L^2(\mathbb{T}^2)$ determined by the matricial symbol function

$$\Psi(z, w) = \begin{bmatrix} \bar{z} - 1 & 0 \\ 0 & \bar{w} - 1 \end{bmatrix}.$$

One can similarly verify the following.

Proposition 5.6. (i) *The matricial symbol function of the kagome framework determined by the six-edge motif is given by*

$$\Phi_{kag}(z, w) = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 1 & -\sqrt{3} & -1 & \sqrt{3} \\ -1 & -\sqrt{3} & 0 & 0 & 1 & \sqrt{3} \\ 2 & 0 & -2z & 0 & 0 & 0 \\ 0 & 0 & -1 & \sqrt{3} & \bar{z}w & -\sqrt{3}\bar{z}w \\ \bar{w} & \sqrt{3}\bar{w} & 0 & 0 & -1 & -\sqrt{3} \end{bmatrix}$$

(ii) *The determinant of $\Phi_{kag}(z, w)$ is a multiple of*

$$zw(\bar{z} - 1)(\bar{w} - 1)(\bar{z} - \bar{w})$$

(iii) *The kagome framework is square-summably isostatic.*

We remark that the function matrix association above has also arisen in engineering in the analysis of Hutchinson and Fleck [19] of the stresses and rigidity of the kagome repetitive truss framework. This is derived from the crystallographic perspective of wave periodic flexes and Bloch's theorem.

5.4. Symmetry equations for infinite frameworks. Let \mathcal{C} be a crystal framework in \mathbb{R}^d with complex Hilbert space rigidity operator R . Also let $\mathcal{K}_v^2, \mathcal{K}_e^2$ be as before, let \mathcal{K}_{fl}^2 be the space of square-summable infinitesimal flexes, let \mathcal{K}_{str}^2 be the space of square-summable infinitesimal stresses and let $\mathcal{M} = \mathcal{K}_v^2 \ominus \mathcal{K}_{fl}^2, \mathcal{N} = \mathcal{K}_e^2 \ominus \mathcal{K}_{str}^2$ be their complementary spaces.

As well as commuting with the coordinate shift operators the rigidity operator satisfies commutation relations for every isometric symmetry of \mathcal{C} . For example, let S be an isometric (not necessarily linear) operator on \mathbb{R}^d which effects a symmetry of \mathcal{C} . There is an induced unitary operator S_e on the complex Hilbert space $\ell^2(E)$ and an analogous operator S_v on $\ell^2(V)$ which permute the standard basis elements. Write \tilde{S}_v for the isometric (not necessarily linear) operator $S_v \otimes S$ on $\mathcal{H}_v = \ell^2(V) \otimes \mathbb{C}^d$ where S is also viewed as an isometric operator on \mathbb{C}^d . Then we have the fundamental symmetry equation $S_e R = R \tilde{S}_v$. (See [27].)

Moreover, if \mathcal{G} is the symmetry group of \mathcal{C} arising from isometric transformations of \mathbb{R}^d and if ρ_e and $\tilde{\rho}_v$ are the associated representations of \mathcal{G} on \mathcal{K}_e^2 and \mathcal{K}_v^2 then we have the symmetry equations

$$\rho_e(g)R = R\tilde{\rho}_v(g) \text{ for } g \in \mathcal{G}.$$

We have shown in [27] how such symmetry equations may be used to obtain a simple proof of a unitary equivalence which implies the Fowler-Guest formula [9] together with various generalisations. In the present setting we have the following analogue which also leads to counting conditions for isostatic and rigid frameworks.

Theorem 5.7. *Let \mathcal{C} be a crystal framework with isometry symmetry group \mathcal{G} and let*

$$\begin{aligned} \tilde{\rho}_v &= \rho_{\mathcal{M}} \oplus \rho_{fl} \\ \rho_e &= \rho_{\mathcal{N}} \oplus \rho_{str} \end{aligned}$$

be the decompositions associated with the spaces of square-summable infinitesimal flexes and stresses. Then $\rho_{\mathcal{M}}$ and $\rho_{\mathcal{N}}$ are unitarily equivalent representations. In particular if \mathcal{C} is square summably isostatic then $\tilde{\rho}_v$ and ρ_e are unitarily equivalent.

Proof. In the square-summably isostatic case R is a bounded operator with trivial kernel and trivial cokernel. The partially isometric part U of the polar decomposition $R = U(R^*R)^{1/2}$ is therefore unitary and it is a standard verification that this unitary also intertwines the representations. In general the symmetry equations show that the space \mathcal{K}_{fl}^2 is reducing for $\tilde{\rho}_v$ and \mathcal{K}_{str}^2 is reducing for ρ_e and so the asserted direct sum decompositions do exist. Now the restriction of R to \mathcal{M} maps to \mathcal{N} with trivial kernel and cokernel and so as before the restriction representations are unitarily equivalent. \square

5.5. From matrix function to wave modes. For a crystal framework $\mathcal{C} = (F_v, F_e, \mathbb{Z}^d)$ with a given motif let $\Phi : \mathbb{T}^d \rightarrow M_{m,nd}$ be the associated matricial symbol function.

Definition 5.8. The mode multiplicity function of \mathcal{C} associated with the given motif and translation group is the function $\mu : \mathbb{T}^d \rightarrow \mathbb{Z}_+$ given by $\mu(z) = \dim \ker \Phi(z)$.

For $d = 2$ we also consider μ as being parametrised by coordinates s, t in $[0, 1) \times [0, 1)$ so that $(z, w) \in \mathbb{T}^2$ corresponds to $(e^{2\pi is}, e^{2\pi it})$. From the determinant calculations above we obtain the following.

Proposition 5.9. (i) For the grid framework $\mathcal{G}_{\mathbb{Z}^2}$ and the 8-edged motif the mode multiplicity function has values $\mu(0, 0) = 2$,

$$\mu(s, 0) = \mu(0, t) = 1,$$

if s and t are nonzero, and is zero otherwise.

(ii) For the kagome framework \mathcal{G}_{kag} and the 6-edged motif the mode multiplicity function has values $\mu(0, 0) = 2$, at the origin while $\mu(s, s) = \mu(s, 0) = \mu(0, t) = 1$ if $s \neq 0$ and $t \neq 0$ and is zero otherwise.

Consider now the *wave flexes* of \mathcal{C} which we define as the infinitesimal flexes which are 1-cell-periodic modulo a phase factor.

Definition 5.10. Let $\mathcal{C} = (F_v, E_v, \mathbb{Z}^d)$ be a crystal framework in \mathbb{R}^d . An (infinitesimal) wave flex of \mathcal{C} is a complex infinitesimal flex $u = (u_v)_{v \in V}$ which is wave periodic (or more precisely 1-cell wave periodic) in the sense that for some vector q in \mathbb{R}^d

$$u_{\kappa+n} = e^{2\pi i \langle q, n \rangle} u_{\kappa}$$

for each vertex κ in the motif set F_v and each d -tuple $n = (n_1, \dots, n_d)$.

The values of the mode multiplicity function corresponds to the dimension of the spaces of wave flexes. To see this suppose that $\Phi(z)$ is the matricial symbol function for $\mathcal{C} = (G, p) = (F_v, F_e, \mathbb{Z}^3)$, that $w \in \mathbb{T}^3$ and that $\det \Phi(w) = 0$. Then $\Phi(w)u_m = 0$ for some nonzero complex motif vector u_m . For the Dirac delta function $\delta_w(z)$ on \mathbb{T}^3 we have, informally, $\Phi(z)F(z) = 0$ for all z on the 3-torus where $F(z)$ is the function $\delta_w(z)u_m$. Thus, taking Fourier transforms it follows that the wave periodic vector $u = \mathcal{F}(F)$ in \mathcal{K}_v satisfies $R(G, p)u = 0$. This shows that the bounded (and non square-summable) vector

$$u = (u_{\kappa, n})_{\kappa \in F_v, n \in \mathbb{Z}^3} = \mathcal{F}(F)(\kappa, n) = w^n u_{\kappa}$$

is a wave flex. The Dirac delta argument can be rigourised in the usual manner and so we obtain the following theorem.

Theorem 5.11. Let $\mathcal{C} = (F_v, F_v, \mathbb{Z}^d)$ be a crystal framework in \mathbb{R}^d with associated symbol function $\Phi(z)$ and suppose that \mathcal{C} has no internal (finitely supported) self-stresses and no internal (finitely supported) flexes. Then

(i) $d|F_v| = |F_e|$, the associated matricial symbol function $\Phi(z)$ on the d -torus is square with full rank almost everywhere and \mathcal{C} is square-summably isostatic.

(ii) infinitesimal wave flexes (that are 1-cell periodic-modulo-phase) exist, with phase factor $w \in \mathbb{T}^d$, if and only if $\det \Phi(w) = 0$. In this case the dimension of the corresponding space of wave flexes is $\dim \ker \Phi(w)$.

It follows also from this that the zero set of the determinant of the matricial symbol function associated with the motif of an appropriate supercell coincides with the phases of supercell-periodic wave flexes.

Of particular computational and theoretical interest is what one might refer to as the wave flex acquisition when a crystal framework deforms under a colossal flex to a framework with higher symmetry. This phenomenon serves as a model for the appearance of so-called Rigid Unit Modes (RUMs) in higher symmetry phases of various material crystals.

The following theorem generalises an interesting result of Wegner [32] for tetrahedral crystals. It may be viewed as an expression of the simple principle that additional symmetry often entails additional flexibility. Our proof applies to arbitrary crystal frameworks and is quite direct, benefitting somewhat from the economy of operator theory formalism. We remark that there are also natural operator algebra perspectives that are relevant to symmetry considerations. The result shows in particular for $d = 3$ that the RUM set is typically a union of surfaces, being the zero set of a single real-valued polynomial, rather than the intersection of the zero sets of the real and imaginary part of a complex polynomial.

For a given crystal framework and motif write Ω for the subset of \mathbb{T}^d formed by the phases ω of the 1-cell periodic wave flexes. We also refer to this informally as the RUM set or the RUMs of \mathcal{C} .

Theorem 5.12. *Let $\mathcal{C} = (F_v, F_e, \mathbb{Z}^d)$ be a crystal framework in \mathbb{R}^d with $d|F_v| = |F_e|$ and suppose that \mathcal{C} possesses inversion symmetry. Then the set Ω has the form*

$$\Omega = \mathbb{T}^d \cap V(p)$$

where $V(p)$ is the zero set of a complex polynomial $p(z_1, \dots, z_d, \bar{z}_1, \dots, \bar{z}_d)$ which is real-valued on \mathbb{T}^d .

Proof. Let us denote the set of framework vertices and edges as

$$\mathcal{V} = \{\kappa + n : \kappa \in F_v, n \in \mathbb{Z}^d\}$$

$$\mathcal{E} = \{e + n : e \in F_e, n \in \mathbb{Z}^d\}$$

Effecting a translation, if necessary, we may assume that the inversion is $\sigma : x \rightarrow -x$, that the unit cell has inversion symmetry and $\sigma(F_v) = F_v$. It may or may not be possible to re-choose the noninternal edges of F_e so that that $\sigma(F_e) = F_e$. Suppose first that this is the case. We show that $\det \Phi(z)$ itself is either real-valued or purely imaginary, from which the stated form for Ω follows.

With the notation of Section 5.4 we have the symmetry equation

$$\rho_e(\sigma)R = R\tilde{\rho}_v(\sigma).$$

Recall that $\rho_e(\sigma)$ and $\tilde{\rho}_v(\sigma)$ are the isometric operators induced by the (isometric) symmetry element σ . Taking Fourier transforms this

equation takes the form

$$U(\sigma)M_{\Phi(z)} = M_{\Phi(z)}V(\sigma)$$

where $U(\sigma)$ and $V(\sigma)$ are the unitary operators determined by their action on the distinguished orthonormal bases. In view of the assumption we have

$$U(\sigma)(\eta_k \otimes z^l) = \eta_{\sigma_e(k)} \otimes z^{-l}, \quad k = 1, \dots, |F_e|$$

(for certain induced permutations σ_e, σ_v) and

$$V(\sigma)(\xi_k^{x_i} \otimes z^l) = \xi_{\sigma_v(k)}^{x_i} \otimes z^{-l}, \quad k = 1, \dots, |F_v|, i = 1, \dots, d.$$

That is, these operators have the form

$$U(\sigma) = E_\sigma \otimes J, \quad V(\sigma) = V_\sigma \otimes J$$

where J is the inversion unitary operator on $L^2(\mathbb{T}^d)$ given by $(Jf)(z) = f(\bar{z})$, and where E_σ and V_σ are scalar permutation matrices induced by σ . Substituting these forms we see that

$$\Phi(z) = (E_\sigma \otimes J)^{-1}\Phi(z)(V_\sigma \otimes J) = E_\sigma^{-1}\Phi(\bar{z})V_\sigma.$$

It follows that

$$\det \Phi(z) = \det \Phi(\bar{z}) \det(E_\sigma)^{-1} \det(V_\sigma) = (-1)^\tau \overline{\det(\Phi(z))}$$

for some integer τ , since the determinant polynomial has real coefficients and so the determinant is either real or purely imaginary.

In the general case because of edges lying on inversion axes (lines through the origin) σ cannot be made to act freely on the motif edges. However for each motif edge index k there is a monomial shift factor $z^{p(k)}$ such that with D the diagonal matrix function

$$D = \text{diag}(z^{p(1)}, \dots, z^{p(d)})$$

we have

$$U(\sigma) = D(E_\sigma \otimes J).$$

Now we see that $\det \Phi(z) = (-1)^\tau z^p \overline{\det(\Phi(z))}$ for some multi-index p . Write $\Phi = F_1 + iF_2$ with the F_i real-valued and $(-1)^\tau z^p = e^{i\gamma(z)}$ with $\gamma = \gamma(z)$ real. Equating real and imaginary parts it follows that either $\cos \gamma = 1$ in which case $\det \Phi(z) = \overline{\det(\Phi(z))}$ and Φ is real-valued, or $F_1(z) = (\sin \gamma)(1 - \cos \gamma)^{-1}F_2(z)$. In all cases, for z on the torus, $F_1(z) = 0$ if and only if $F_2(z) = 0$, as required. \square

5.6. Honeycomb frameworks and the kagome net. The explicit formulation of the matricial symbol function makes clear a simple additive principle which is useful for calculation : adding internal edges to a unit cell results in adding the same number of rows to the symbol function to form the new symbol function, and each of these rows has the standard form with constant entries.

Consider the following honeycomb frameworks based on regular hexagons.




FIGURE 19. A 3-regular honeycomb framework.




FIGURE 20. A 4-regular honeycomb framework.

The 3-regular (symmetric) honeycomb framework has a 9 by 12 matricial symbol function. There are vectors in the kernel of the multiplication operator providing internal infinitesimal flexes and there are evident bounded infinitesimal flexes associated with "parallel deformations", for example.

A 4-regular framework may be constructed by rigidifying the hexagon internal to the unit cell, by adding three cross edges. The matricial symbol function is correspondingly enlarged by three scalar rows, appearing in rows 9 to 12 in the following matrix.

$$\begin{bmatrix}
 -1/3 & 0 & 1/3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & -1/6 & -1/6\sqrt{3} & 1/6 & 1/6\sqrt{3} & 0 & 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1/6 & -1/6\sqrt{3} & -1/6 & 1/6\sqrt{3} & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 1/3 & 0 & -1/3 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1/6 & 1/6\sqrt{3} & -1/6 & -1/6\sqrt{3} \\
 1/6 & -1/6\sqrt{3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1/6 & 1/6\sqrt{3} \\
 0 & 0 & 0 & 0 & -1/3 & 0 & 0 & 0 & 0 & 0 & 1/3\bar{z} & 0 \\
 1/6\bar{w} & 1/6\sqrt{3}\bar{w} & 0 & 0 & 0 & 0 & -1/6 & -1/6\sqrt{3} & 0 & 0 & 0 & 0 \\
 0 & 0 & -1/6\bar{w} & 1/6\sqrt{3}\bar{w} & 0 & 0 & 0 & 0 & 1/6\bar{z} & -1/6\sqrt{3}\bar{z} & 0 & 0 \\
 -1/3 & -1/3\sqrt{3} & 0 & 0 & 0 & 0 & 1/3 & 1/3\sqrt{3} & 0 & 0 & 0 & 0 \\
 0 & 0 & 1/3 & -1/3\sqrt{3} & 0 & 0 & 0 & 0 & -1/3 & 1/3\sqrt{3} & 0 & 0 \\
 0 & 0 & 0 & 0 & 2/3 & 0 & 0 & 0 & 0 & 0 & -2/3 & 0
 \end{bmatrix}$$

The cross-barred hexagon is continuously rigid but is infinitesimally flexible. We refer to this flex as the in-out flex. (See Whiteley [34] for related discussions.) It is straightforward to show that this flex extends periodically as a 1-cell-periodic infinitesimal flex of the framework.

The motif matrix function is obtained by evaluating at $z = w = 1$. This has rank 8 reflecting the four independent 1-cell-periodic flexes corresponding to horizontal translation, to vertical translation, to local rotation of the cross-barred hexagon, and to the periodic "in-out" flex.

We define the kagome net framework as the three dimensional crystal framework $\mathcal{G}_{knet} = (F_v, F_e, \mathbb{Z}^3)$ determined by the motif diagram in Figure 21. This figure shows a tetrahedron internal to a parallelepiped unit cell together with six additional non-internal edges, each of which extends a tetrahedron edge as indicated.




FIGURE 21. A twelve-edged motif for the kagome net, \mathcal{G}_{knet} .

For the given motif the motif rigidity matrix $R_m(\mathcal{G}_{knet})$ takes the form of a 12 by 12 matrix valued function on the 3-torus. Different motifs, for the same translation group, give matrix functions that are related by permutations on columns and monomial multiplication on rows. It can be shown that the determinant of such a matrix function

is equal to a scalar and monomial multiple of the polynomial

$$(z - 1)(w - 1)(u - 1)(z - w)(w - u)(u - z).$$

This is in complete analogy with the kagome framework and in the light of earlier discussions may draw the following conclusions.

Theorem 5.13. *(i) The kagome net \mathcal{G}_{knet} is square-summably isostatic and possesses no internal infinitesimal flexes.*

(ii) The mode multiplicity function $\mu : \mathbb{T}^3 \rightarrow \mathbb{Z}_+$ of the kagome net framework has support equal to the intersection of \mathbb{T}^3 with the six planes

$$z = 1, w = 1, u = 1, z = w, w = u, u = z.$$

5.7. Crystallography and Rigid Unit Modes. Rigidity theory and a Hilbert space operator viewpoint have led us to determine, ab initio, the matricial symbol function $\Phi(z)$ of an abstract crystal framework and motif. The zeros of $\det \Phi(z)$ correspond to the phases of infinitesimal periodic-modulo-phase wave flexes and the mode multiplicity function $\mu(w)$ of Φ detects the multiplicities of independent wave flexes for w . There are close connections between these mathematical observations and the Rigid Unit Modes (RUMs) that are observed in certain material crystals through diffraction experiments. This connection is part of the motivation for the formulation of the function $\mu(z)$. It seems to us that the explicit algorithm for the passage from crystal motif to matrix function will provide a useful computational and theoretical tool for identifying RUMs and their relationships.

Infinitesimal wave flexes for abstract crystal frameworks appear in the classical crystallography of Born von Karman theory. For example in the case of variants of quartz the low energy excitation modes result from rigid unit motions of SiO_4 molecular tetrahedra within a tetrahedral net. These modes are therefore modelled by the first order periodic-modulo-phase infinitesimal rigid motions of abstract tetrahedral net frameworks. For this connection see, for example, Chapter 3 of [35]. Comparisons of extensive simulations and experimental results have shown that the RUM modes of simulated crystals are closely correlated with observed RUMs. For this see the seminal paper of Giddy et al [11], and also, for example, Dove et al [8] and Goodwin et al [14].

An interesting advance has been obtained recently by Wegner [32] who has derived mathematically, rather than through simulation, the RUM sets for various idealised tetrahedral crystals that model the geometry of material crystals, including β -cristobalite, HP tridymite, β -quartz, α -cristobalite and α -quartz. These models are standard framework models that coincide with crystal bar-joint frameworks in our terminology and the results are obtained by determining vanishing determinants (zero sets of $\det \Phi$ in our formalism) by means of computer algebra and symmetry reductions. In particular Wegner obtains an

analytic derivation of some of the curious surfaces observed in experiments of Dove et al.

REFERENCES

- [1] L. Asimow and B. Roth, The rigidity of graphs, *Trans. Amer. Math. Soc.*, 245 (1978), 279-289.
- [2] C.S. Borcea and I. Streinu, Periodic frameworks and flexibility, *Proc. R. Soc. A*, doi:10.1098/rspa.2009.0676.
- [3] A. Böttcher, B. Silberman, *Analysis of Toeplitz operators*, Springer-Verlag, Berlin, 1990.
- [4] R. Connelly, P.W. Fowler, S.D. Guest, B. Schulze, W.J. Whiteley, "When is a pin-jointed framework isostatic?", *International Journal of Solids and Structures*, 46 (2009), 762 - 773.
- [5] D. Cox, J. Little, D. O'Shea, *Ideals, Varieties and Algorithms*, Springer-Verlag, 1992.
- [6] M. V. Chubynsky and M. F. Thorpe, Algorithms for three-dimensional rigidity analysis and a first-order percolation transition, *Physical Review E* 76, 2007.
- [7] A. Donev and S. Torquato, Energy-efficient actuation in infinite lattice structures, *J. Mech Phys. Solids*, 51 (2003) 1459-1475.
- [8] M. T. Dove et al, Floppy Modes in Crystalline and Amorphous Silicates; in *Rigidity Theory and Applications*, Edited by M. F. Thorpe and P. M. Duxbury, Kluwer Academic/Plenum Publishers, New York (1999).
- [9] P.W. Fowler and S.D. Guest, A symmetry extension of Maxwell's rule for rigidity of frames, *International Journal of Solids and Structures* 37 (2000) 1793 - 1804.
- [10] M. G. Frost and C. Storey, Equivalence of a matrix over $\mathcal{R}[s, z]$ with its Smith form, *Int. J. Control*, 28 (1978), 665-671.
- [11] A. P. Giddy, M.T. Dove, G.S. Pawley, V. Heine, The determination of rigid unit modes as potential soft modes for displacive phase transitions in framework crystal structures. *Acta Crystallogr.*, A49 (1993), 697 - 703.
- [12] H. Gluck, Almost all simply connected closed surfaces are rigid, in *Geometric Topology*, Lecture Notes in Math., no. 438, Springer-Verlag, Berlin, 1975, pp. 225-239.
- [13] I. Gohberg, P. Lancaster, L. Rodman, *Matrix Polynomials*, Classics in Applied Mathematics, SIAM, 2009.
- [14] A. L. Goodwin, M. Calleja, M. J. Conterio, M. T. Dove, J. S. O. Evans, D. A. Keen, L. Peters, M. G. Tucker, Colossal Positive and Negative Thermal Expansion in the Framework Material $\text{Ag}_3[\text{Co}(\text{CN})_6]$, *Science* 8 February 2008: Vol. 319. no. 5864, pp. 794 - 797.
- [15] J. Graver, B. Servatius and H. Servatius, *Combinatorial rigidity*, Graduate Texts in Mathematics, vol 2, Amer. Math. Soc., 1993.
- [16] S. D. Guest and J. W. Hutchinson, On the determinacy of repetitive structures, *Journal of the Mechanics and Physics of Solids*, 51 (2003), 383 - 391.
- [17] B. Grunbaum, G.C. Shephard, *Tilings and patterns*, W.H. Freeman, New York, 1987.
- [18] K.D. Hammonds, M.T. Dove, A.P. Giddy, V. Heine, B. Winkler, Rigid-unit phonon modes and structural phase transitions in framework silicates, *American Mineralogist*, 81 (1996), 1057-1079.
- [19] R.G. Hutchinson and N.A. Fleck, The structural performance of the periodic truss, *Journal of the Mechanics and Physics of Solids*, 54 (2006) 756-782.
- [20] A. B. Kempe, On a general method of describing plane curves of the nth degree by linkwork, *Proc. London Math. Soc.*, 7 (1876), 213-216.

- [21] G. Laman, On graphs and the rigidity of plane skeletal structures, *J. Engineering Mathematics*, 4 (1970), 331-340.
- [22] S. Lang, *Algebra*, Addison Wesley, 1993.
- [23] J.C. Maxwell, On the calculation of the equilibrium and stiffness of frames. *Philosophical Magazine* 27, (1864), 294 - 299.
- [24] J.C. Owen and S.C. Power, The non-solvability by radicals of generic 3-connected planar Laman graphs, *Trans. Amer. Math. Soc.*, 359 (2007), 2269-2303.
- [25] J.C. Owen and S.C. Power, Infinite bar-joint frameworks, *Proceedings of the Symposium in Applied Computing, (SAC 2009) March 8-12, 2009, Honolulu, Hawaii, U.S.A.*
- [26] J.C. Owen and S.C. Power, Continuous curves from infinite Kempe linkages, *Bull. London Math. Soc.*, 2009; doi: 10.1112/blms/bdp087
- [27] J.C. Owen and S.C. Power, Frameworks, symmetry and rigidity, to appear in the *Inter. J. of Computational Geometry and Applications*, arXiv:0812.3785.
- [28] E. Ross, private communication, July 2009.
- [29] B. Roth, Rigid and flexible frameworks, *American Math. Monthly*, Vol. 88 (1981), 6-21.
- [30] W. Rudin, *Functional Analysis*, McGraw and Hill, NY, 1991.
- [31] M. F. Thorpe and P. M. Duxbury, *Rigidity Theory and Applications*, Edited by Kluwer Academic/Plenum Publishers, New York (1999).
- [32] F. Wegner, Rigid-unit modes in tetrahedral crystals, *Journal of Physics: Condensed Matter*, Volume 19, Issue 40, (2007).
- [33] S.A. Wells, M. T. Dove, M. G. Tucker and K. Trachenko, Real-space rigid-unit-mode analysis of dynamic disorder in quartz, cristobalite and amorphous silica, *J. Phys.: Condens. Matter* 14 (2002), 46454657.
- [34] W. Whiteley, Matroids and rigid structures, in "Matroid Applications" ed. N. White, *Encyclopedia of Mathematics and its applications* 40 (1992), 1-51.
- [35] B.T.M. Willis and A.W. Pryor, *Thermal vibrations in crystallography*, Cambridge University Press, 1975.
- [36] M. Wyart, Rigidity-based approach to the boson peak in amorphous solids: from sphere packing to amorphous silica, preprint 2008, arXiv:0806.4596v1.

D-CUBED, SIEMENS PLM SOFTWARE, PARK HOUSE,, CASTLE PARK, CAMBRIDGE UK

E-mail address: owen.john.ext@siemens.com

DEPT. MATH. STATS., LANCASTER UNIVERSITY, LANCASTER LA1 4YF, U.K.

E-mail address: s.power@lancaster.ac.uk