

The real exponential integrals

Notes by G.J.O. Jameson

The functions $E(x)$ and $E^*(x)$

Define, for suitable x ,

$$E(x) = \int_x^\infty \frac{e^{-t}}{t} dt, \quad E^*(x) = \int_0^x \frac{1 - e^{-t}}{t} dt. \quad (1)$$

$E(x)$, as well as various mutations and equivalent forms, is known as the “incomplete exponential integral”. It is the special case $\Gamma(0, x)$ of the “incomplete gamma function”, defined by $\Gamma(a, x) = \int_x^\infty t^{a-1} e^{-t} dt$ (e.g. see [Jam3]). Notation sometimes used in specialist literature is $E_1(x)$ for our $E(x)$ and $\text{Ein}(x)$ for $E^*(x)$.

Clearly, $E(x)$ is defined for all $x > 0$, but not for $x = 0$, since $\int_0^1 e^{-t}/t dt$ is divergent. Since the integrand is positive, $E(x)$ is non-negative and decreasing. By the fundamental theorem of calculus, $E'(x) = -e^{-x}/x$. A simple inequality for $E(x)$ is

$$E(x) \leq \frac{1}{x} \int_x^\infty e^{-t} dt = \frac{e^{-x}}{x}. \quad (2)$$

In particular, $E(x)$, and even $e^x E(x)$, tends to 0 as $x \rightarrow \infty$.

There is no problem about convergence of $E^*(x)$ at 0. In fact, $0 < 1 - e^{-t} \leq t$ for $t > 0$, so $0 < (1 - e^{-t})/t < 1$, and hence $0 < E^*(x) \leq x$ for $x > 0$. In particular, $E^*(x) \rightarrow 0$ as $x \rightarrow 0^+$. A better inequality for $x > 1$ is

$$E^*(x) < E^*(1) + \int_1^x \frac{1}{t} dt < \log x + 1. \quad (3)$$

Also, $E^*(x)$ is increasing and $(E^*)'(x) = (1 - e^{-x})/x$.

By integrating the series

$$\frac{1 - e^{-t}}{t} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{t^{n-1}}{n!},$$

we obtain a power series expression for $E^*(x)$:

$$E^*(x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n!n} = x - \frac{x^2}{2!2} + \frac{x^3}{3!3} - \cdots, \quad (4)$$

from which, in principle, values can be calculated (though in practice the calculation is only pleasant for fairly small x). For example, eight terms are enough to give $E^*(1) \approx 0.79660$ to five decimal places.

We mention some equivalent forms given by substitutions. For $a > 0$, the substitution $at = u$ gives

$$\int_x^\infty \frac{e^{-at}}{t} dt = \int_{ax}^\infty \frac{e^{-u}}{u} du = E(ax), \quad (5)$$

$$\int_0^x \frac{1 - e^{-at}}{t} dt = \int_0^{ax} \frac{1 - e^{-u}}{u} du = E^*(ax), \quad (6)$$

In particular, taking $x = 1$, we obtain

$$E(a) = \int_1^\infty \frac{e^{-at}}{t} dt, \quad E^*(a) = \int_0^1 \frac{1 - e^{-at}}{t} dt. \quad (7)$$

The substitution $t = x + u$ in (1) gives

$$E(x) = e^{-x} \int_0^\infty \frac{e^{-u}}{x + u} du. \quad (8)$$

This expression can be used to define $E(z)$ for any complex z not on the negative real line. As a contour integral, it equates to the integral of $e^{-\zeta}/\zeta$ on the horizontal half-line defined by $\zeta = z + u$ for $u \geq 0$. Not very helpfully, the notation $\text{Ei}(z)$ is sometimes used for $-E(-z)$. We will not discuss the complex function $E(z)$ in these notes.

Substituting $e^{-u} = v$ in (8), we obtain

$$E(x) = e^{-x} \int_0^1 \frac{1}{x - \log v} dv. \quad (9)$$

The numbers $E(1)$ and $E^*(1)$ have particular significance. Integration by parts gives

$$E(1) = \left[e^{-t} \log t \right]_1^\infty + \int_1^\infty e^{-t} \log t dt = \int_1^\infty e^{-t} \log t dt. \quad (10)$$

Since $(1 - e^{-t}) \log t \rightarrow 0$ as $t \rightarrow 0^+$, we have similarly

$$E^*(1) = \left[(1 - e^{-t}) \log t \right]_0^1 - \int_0^1 e^{-t} \log t dt = - \int_0^1 e^{-t} \log t dt. \quad (11)$$

To relate $E^*(x)$ and $E(x)$, observe that

$$E^*(x) - E^*(1) = \int_1^x \frac{1 - e^{-t}}{t} dt = \log x - \int_1^x \frac{e^{-t}}{t} dt = \log x - E(1) + E(x),$$

so

$$E(x) = E^*(x) - \log x + c, \quad (12)$$

where c is constant, in fact $c = E(1) - E^*(1)$. Furthermore, we have two ways to express c as a limit. Since $\lim_{x \rightarrow 0^+} E^*(x) = 0$, we have

$$c = \lim_{x \rightarrow 0^+} [E(x) + \log x] \quad (13)$$

and since $\lim_{x \rightarrow \infty} E(x) = 0$, we have

$$-c = \lim_{x \rightarrow \infty} [E^*(x) - \log x]. \quad (14)$$

Also, by (10) and (11), we have

$$c = \int_0^\infty e^{-t} \log t \, dt. \quad (15)$$

This is known as the (complete) exponential integral (as already noted, e^{-t}/t does not give a convergent integral on $(0, \infty)$). Readers familiar with the gamma function will recognise that it equates to $\Gamma'(1)$.

The evaluation of c will be our main theorem, and we turn to it shortly. But there are some facts that can be deduced without knowing the value of c . Firstly, $E(x) \sim -\log x$ as $x \rightarrow 0^+$, hence $x E(x) \rightarrow 0$ as $x \rightarrow 0^+$. Another deduction is the following integral:

PROPOSITION 1. For $a, b > 0$,

$$\int_0^\infty \frac{e^{-at} - e^{-bt}}{t} \, dt = \log b - \log a. \quad (16)$$

Proof. By (6) and (12),

$$\begin{aligned} \int_0^x \frac{e^{-at} - e^{-bt}}{t} \, dt &= E^*(bx) - E^*(ax) \\ &= E(bx) - E(ax) + \log bx - \log ax \\ &= E(bx) - E(ax) + \log b - \log a \\ &\rightarrow \log b - \log a \quad \text{as } x \rightarrow \infty. \quad \square \end{aligned}$$

A well-known alternative proof of (16) is by expressing the integrand as $\int_a^b e^{-ty} \, dy$ and reversing the double integral. It is also a special case of the ‘‘Frullani integral’’ (see [Fer, p. 133–135]).

Determination of c ; the complete exponential integral

We will use (14) to evaluate c .

LEMMA 1. For $0 \leq t \leq n$,

$$\left(1 - \frac{t^2}{n}\right) e^{-t} \leq \left(1 - \frac{t}{n}\right)^n \leq e^{-t}. \quad (17)$$

Proof. From the series for e^x and $1/(1-x)$, we have $1+x \leq e^x \leq 1/(1-x)$, and hence $(1-x^2)e^{-x} \leq 1-x \leq e^{-x}$ for $0 < x < 1$. Substitute $x = t/n$ and take the n th power

to obtain

$$\left(1 - \frac{t^2}{n^2}\right)^n e^{-t} \leq \left(1 - \frac{t}{n}\right)^n \leq e^{-t}.$$

for $0 \leq t \leq n$. Now $(1 - a)^n \geq 1 - na$ for $0 \leq a \leq 1$, so (17) follows. \square

THEOREM 2. *We have $c = -\gamma$, where γ is Euler's constant. Hence*

$$\int_0^\infty e^{-t} \log t \, dt = -\gamma, \quad (18)$$

$$E^*(x) = E(x) + \log x + \gamma. \quad (19)$$

Proof. By (14), it is enough to show that $E^*(n) - \log n \rightarrow \gamma$ as $n \rightarrow \infty$. Since $e^{-t} = \lim_{n \rightarrow \infty} (1 - \frac{t}{n})^n$, it seems at least plausible that $E^*(n)$ is approximated by K_n , where

$$\begin{aligned} K_n &= \int_0^n \frac{1}{t} \left[1 - \left(1 - \frac{t}{n}\right)^n \right] dt \\ &= \int_0^1 \frac{1}{u} [1 - (1 - u)^n] du \\ &= \int_0^1 \frac{1 - v^n}{1 - v} dv \\ &= \int_0^1 (1 + v + \cdots + v^{n-1}) dv \\ &= 1 + \frac{1}{2} + \cdots + \frac{1}{n}, \end{aligned}$$

so that $K_n - \log n \rightarrow \gamma$ as $n \rightarrow \infty$.

To make this precise, we have by (17)

$$1 - e^{-t} \leq 1 - \left(1 - \frac{t}{n}\right)^n \leq 1 - e^{-t} + \frac{1}{n} t^2 e^{-t}.$$

It follows that $E^*(n) \leq K_n \leq E^*(n) + \Delta_n$, where

$$\Delta_n = \frac{1}{n} \int_0^n t e^{-t} dt < \frac{1}{n}. \quad \square$$

COROLLARY 2.1. *For all $x > 0$,*

$$-\log x - \gamma < E(x) < -\log x - \gamma + x, \quad (20)$$

$$\log x + \gamma < E^*(x) < \log x + \gamma + \frac{e^{-x}}{x}. \quad \square \quad (21)$$

Of course, (20) is effective for small x , and (21) for large x .

Also, we can deduce particular values of $E(x)$, for example $E(1) = E^*(1) - \gamma \approx 0.21938$.

The integral (18), together with most of the equivalent versions described below, was discovered by Euler. The proof given here follows [WW, p. 242]. A slight variant is given in [Lo]. We now record some integrals derived from it, or equivalent to it.

First, the substitution $at = u$ (where $a > 0$) gives

$$\int_0^\infty e^{-at} \log t \, dt = \int_0^\infty e^{-u} (\log u - \log a) \frac{1}{a} \, du = -\frac{1}{a}(\gamma + \log a), \quad (22)$$

which we can rewrite rather pleasantly as $\int_0^\infty e^{-at} (\log t + \gamma) \, dt = -\frac{1}{a} \log a$.

Next, substituting in turn $e^{-t} = u$ and $e^t = u$, we obtain

$$\int_0^1 \log(-\log u) \, du = \int_1^\infty \frac{\log \log u}{u^2} \, du = -\gamma. \quad (23)$$

PROPOSITION 3. *We have*

$$\int_0^\infty t e^{-t} \log t \, dt = 1 - \gamma \quad (24)$$

(Remark: this integral equates to $\Gamma'(2)$.)

Proof. Denote the integral by I . Expressing the integrand as $e^{-t}(t \log t)$ and integrating by parts, we obtain

$$I = \left[-e^{-t}(t \log t) \right]_0^\infty + \int_0^\infty e^{-t} (\log t + 1) \, dt = 0 - \gamma + 1. \quad \square$$

In a similar way, one can prove by induction that

$$\int_0^\infty t^n e^{-t} \log t \, dt = n!(H_n - \gamma),$$

where $H_n = \sum_{r=1}^n \frac{1}{r}$. (Hence $\psi(n+1) = H_n - \gamma$, where $\psi(x) = \Gamma'(x)/\Gamma(x)$.)

We now derive some further integral expressions for γ . Now that we know that $c = -\gamma$, we can apply (14) again to obtain at once:

PROPOSITION 4. *We have*

$$\gamma = \int_0^\infty \frac{1}{t} \left(\frac{1}{t+1} - e^{-t} \right) dt. \quad (25)$$

Proof. By (14),

$$\begin{aligned} \gamma = \lim_{x \rightarrow \infty} [E^*(x) - \log(x+1)] &= \lim_{x \rightarrow \infty} \int_0^x \left(\frac{1 - e^{-t}}{t} - \frac{1}{t+1} \right) dt \\ &= \int_0^\infty \left(\frac{1}{t(t+1)} - \frac{e^{-t}}{t} \right) dt. \quad \square \end{aligned}$$

Alternatively, (25) can be derived from (18) by using (16) to substitute for $\log x$ and reversing the resulting double integral.

PROPOSITION 5. *We have*

$$\gamma = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{e^{-t}}{t} \right) dt. \quad (26)$$

Proof. This will follow from (25) if we prove that

$$\int_0^\infty \left(\frac{1}{t(t+1)} - \frac{1}{e^t - 1} \right) dt = 0.$$

For $\delta > 0$, denote by I_δ this integral reduced to the interval $[\delta, \infty)$. Noting that $1/(e^t - 1) = e^{-t}/(1 - e^{-t})$, we have

$$I_\delta = \log(1 + \delta) - \log \delta + \log(1 - e^{-\delta}).$$

Now $\frac{1}{\delta}(1 - e^{-\delta}) \rightarrow 1$ as $\delta \rightarrow 0^+$, hence $I = \lim_{\delta \rightarrow 0^+} I_\delta = 0$. \square

Alternatively, one can prove (26) directly and derive (18) [BM, Proposition 9.3.3]. The substitution $e^{-t} = x$ in (26) gives:

COROLLARY 5.1. *We have*

$$\gamma = \int_0^1 \left(\frac{1}{1-x} + \frac{1}{\log x} \right) dx. \quad \square \quad (27)$$

(25) and (26) are the cases $x = 1$ in the two standard integral expressions for $\psi(x) = \Gamma'(x)/\Gamma(x)$ [AAR, p. 26].

PROPOSITION 6. [Sco] *We have*

$$\gamma = 2 \int_0^\infty \frac{1}{t} (e^{-t^2} - e^{-t}) dt. \quad (28)$$

Proof. Recall that $\gamma = E^*(1) - E(1)$. Substituting $t = u^2$, we have

$$E(1) = \int_1^\infty \frac{e^{-t}}{t} dt = 2 \int_1^\infty \frac{e^{-u^2}}{u} du,$$

hence

$$E(1) = 2E(1) - E(1) = 2 \int_1^\infty \frac{1}{t} (e^{-t} - e^{-t^2}) dt.$$

Similarly,

$$E^*(1) = \int_0^1 \frac{1 - e^{-t}}{t} dt = 2 \int_0^1 \frac{1 - e^{-u^2}}{u} du,$$

so

$$E^*(1) = 2E^*(1) - E^*(1) = 2 \int_0^1 \frac{1}{t}(e^{-t^2} - e^{-t}) dt.$$

Statement (28) follows. \square

Calculation of γ . The identity $\gamma = E^*(x) - E(x) - \log x$ has been used for the calculation of γ to great degrees of accuracy. For a suitably chosen x , one can calculate $E^*(x)$ and $\log x$, and estimate $E(x)$ by its asymptotic expansion (which we discuss next), or simply choose x so that $E(x)$ is suitably small. For a survey of this topic, see [GS], [Lag] or [BM, chap. 9].

Asymptotic expression for $E(x)$

The inequality (2) is enough for many purposes, but there is an asymptotic expression giving a better estimation for large x , if wanted. It is helpful to describe the process more generally, as follows. Suppose that $f(t)$ and all its derivatives tend to 0 as $t \rightarrow \infty$, and write $I_f(x) = \int_x^\infty e^{-t} f(t) dt$. Integrating by parts, we have

$$I_f(x) = \left[-e^{-t} f(t) \right]_x^\infty + \int_x^\infty e^{-t} f'(t) dt = e^{-x} f(x) + I_{f'}(x).$$

Applying this to $f'(t)$ and substituting, and repeating the process, we obtain for all $k \geq 1$

$$I_f(x) = e^{-x} [f(x) + f'(x) + \cdots + f^{(k-1)}(x)] + I_{f^{(k)}}(x).$$

If also $(-1)^k f^{(k)}(t)$ is non-negative and decreasing, then $0 \leq (-1)^k I_{f^{(k)}}(x) \leq (-1)^k e^{-x} f^{(k)}(x)$. For $E(x)$, we apply this with $f(x) = 1/x$. The conditions are satisfied, so we obtain

$$E(x) = e^{-x} \left(\frac{1}{x} - \frac{1}{x^2} + \frac{2}{x^3} - \cdots + (-1)^{k-1} \frac{(k-1)!}{x^k} + (-1)^k r_k(x) \right), \quad (29)$$

where

$$0 \leq r_k(x) \leq \frac{k!}{x^{k+1}}.$$

For large x , we can derive a fairly accurate estimation of $E(x)$ by choosing a suitable number of terms. For example $E(10) = \rho e^{-10}$, where $0.0915 < \rho < 0.0917$. However, what this generates is an *asymptotic expression*, not a convergent series, because for any particular x , the successive derivatives ultimately grow large.

Some integrals involving $E(x)$

We record some integrals involving $E(x)$. None of them require the fact that $c = -\gamma$. For the first one, recall that $\int_0^\infty t^{a-1} e^{-t} dt$ defines $\Gamma(a)$ for $a > 0$. We compare two methods.

PROPOSITION 7. For $a > 0$,

$$\int_0^\infty x^{a-1} E(x) dx = \frac{\Gamma(a)}{a}. \quad (30)$$

In particular,

$$\int_0^\infty E(x) dx = 1. \quad (31)$$

Proof 1. Reversing the double integral, we obtain

$$\begin{aligned} \int_0^\infty x^{a-1} E(x) dx &= \int_0^\infty x^{a-1} \int_x^\infty \frac{e^{-t}}{t} dt dx \\ &= \int_0^\infty \frac{e^{-t}}{t} \int_0^t x^{a-1} dx dt \\ &= \frac{1}{a} \int_0^\infty t^{a-1} e^{-t} dt \\ &= \frac{\Gamma(a)}{a}. \end{aligned}$$

Proof 2. Integrate by parts:

$$\int_0^\infty x^{a-1} E(x) dx = \left[\frac{x^a}{a} E(x) \right]_0^\infty + \int_0^\infty \frac{x^a}{a} \frac{e^{-x}}{x} dx.$$

Now $x^a E(x) \rightarrow 0$ as $x \rightarrow \infty$. Also, since $E(x) \sim -\log x$ for x close to 0, $x^a E(x) \rightarrow 0$ as $x \rightarrow 0$. So the integral equates to $\frac{1}{a} \int_0^\infty x^{a-1} e^{-x} dx = \Gamma(a)/a$. \square

So for positive integers n , $\int_0^\infty x^n E(x) dx = n!/(n+1)$.

PROPOSITION 8. For $a > 0$,

$$\int_0^\infty e^{-ax} E(x) dx = \frac{1}{a} \log(a+1). \quad (32)$$

Proof. The integral is

$$\int_0^\infty e^{-ax} \int_x^\infty \frac{e^{-t}}{t} dt dx = \int_0^\infty \frac{e^{-t}}{t} \int_0^t e^{-ax} dx dt = \int_0^\infty \frac{e^{-t}(1 - e^{-at})}{at} dt.$$

The stated value now follows, by (16). \square

PROPOSITION 9. We have

$$\int_0^\infty E(x)^2 dx = 2 \log 2. \quad (33)$$

Proof. Integrate by parts:

$$\begin{aligned} \int_0^\infty 1 \cdot E(x)^2 dx &= \left[xE(x)^2 \right]_0^\infty + 2 \int_0^\infty xE(x) \frac{e^{-x}}{x} dx \\ &= 2 \int_0^\infty e^{-x} E(x) dx \\ &= 2 \log 2, \end{aligned}$$

in which we used (32) and the fact that $\lim_{x \rightarrow 0^+} [xE(x)^2] = 0$. \square

To prove (33) by the double-integral method, one would start by expressing $E(x)^2$ as an integral:

$$E(x)^2 = - \int_x^\infty 2E(t)E'(t) dt = 2 \int_x^\infty E(t) \frac{e^{-t}}{t} dt.$$

Application to the cosine integral

Consider the integrals corresponding to (1) with e^{-t} replaced by $\cos t$:

$$C(x) = \int_x^\infty \frac{\cos t}{t} dt, \quad C^*(x) = \int_0^x \frac{1 - \cos t}{t} dt. \quad (34)$$

Convergence of the integral defining $C(x)$, together with the fact that $\lim_{x \rightarrow \infty} C(x) = 0$, is easily established by integration by parts. We also define (following standard notation)

$$\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt.$$

We will use a contour integral to relate $C^*(x)$ and $E^*(x)$, enabling us derive a result analogous to Theorem 2 for $C(x)$.

Exactly as for $E(x)$, we have

$$C(x) = C^*(x) - \log x + c', \quad (35)$$

where $c' = C(1) - C^*(1)$. By (12) and (35),

$$C(x) - E(x) = C^*(x) - E^*(x) + c' - c. \quad (36)$$

We will prove:

THEOREM 10. *We have $C^*(x) - E^*(x) \rightarrow 0$ as $x \rightarrow \infty$, hence $c' = -\gamma$ and*

$$C(x) = C^*(x) - \log x - \gamma, \quad (37)$$

$$\int_0^\infty \frac{\cos t - e^{-t}}{t} dt = 0. \quad (38)$$

Once we have shown that $C^*(x) - E^*(x) \rightarrow 0$ as $x \rightarrow \infty$, it follows from (36) that $c' = c$, so $c' = -\gamma$, hence (37). Also, it follows that $C(x) - E(x) \rightarrow 0$ as $x \rightarrow 0^+$, hence (38) (for this, we do not need to know that $c = -\gamma$).

LEMMA 2. *We have*

$$0 \leq \int_0^{\pi/2} e^{-R \sin \theta} d\theta \leq \frac{\pi}{2R}.$$

Proof. Since the function $\sin \theta$ is concave on $[0, \frac{\pi}{2}]$, we have $\sin \theta \geq \frac{2\theta}{\pi}$ on $[0, \frac{\pi}{2}]$. Hence

$$\int_0^{\pi/2} e^{-R \sin \theta} d\theta \leq \int_0^{\pi/2} e^{-2R\theta/\pi} d\theta = \left[-\frac{\pi}{2R} e^{-2R\theta/\pi} \right]_0^{\pi/2} = \frac{\pi}{2R} (1 - e^{-R}). \quad \square$$

Proof of Theorem 10. Let C_R be the circular arc of radius R in the positive quadrant, represented by $z = Re^{i\theta}$ for $0 \leq \theta \leq \frac{\pi}{2}$. Denote by Γ the closed contour consisting of C_R together with the real interval $[0, R]$, and the imaginary axis from iR to 0 . Let

$$f(z) = \frac{1 - e^{iz}}{z}.$$

Then f has no pole at 0 , since $f(z) = -i + \frac{1}{2}z + \dots$. By Cauchy's integral theorem, $\int_{\Gamma} f(z) dz = 0$. The contribution of the real axis is

$$\int_0^R \frac{1 - e^{it}}{t} dt = C^*(R) - i \operatorname{Si}(R).$$

The contribution of the imaginary axis, taken towards the origin, is

$$- \int_0^R \frac{1 - e^{-t}}{t} dt = -E^*(R).$$

Now consider C_R . The contribution of the term $\frac{1}{z}$ is $\frac{\pi}{2}i$. The contribution of the other term is

$$I_R =: \int_{C_R} \frac{e^{iz}}{z} dz = \int_0^{\pi/2} i e^{iRe^{i\theta}} d\theta.$$

Now $|e^{iRe^{i\theta}}| = e^{-R \sin \theta}$, so by Lemma 2, $|I_R| \leq \pi/(2R)$.

Now considering the real part, we deduce

$$|C^*(R) - E^*(R)| \leq \frac{\pi}{2R},$$

so $C^*(R) - E^*(R) \rightarrow 0$ as $R \rightarrow \infty$. \square

At the same time, we deduce from the imaginary part that $|\operatorname{Si}(R) - \frac{\pi}{2}| \leq \frac{\pi}{2R}$, thereby proving the ‘‘sine integral’’

$$\int_0^{\infty} \frac{\sin t}{t} dt = \frac{\pi}{2}.$$

An alternative proof of Theorem 10 by double integrals is given in [Jam2]. A direct proof of (37) can be seen in [Jam1].

References

- [AAR] George E. Andrews, Richard Askey and Ranjan Roy, *Special Functions*, Cambridge Univ. Press (1999).
- [BM] George Boros and Victor H. Moll, *Irresistible Integrals*, Cambridge Univ. Press (2004).
- [Fer] W. L. Ferrar, *Integral Calculus*, Oxford Univ. Press (1958).
- [GS] Xavier Gourdon and Pascal Sebah, The Euler constant γ , at <http://numbers.computation.free.fr/Constants/constants.html>.
- [Jam1] G. J. O. Jameson, Sine, cosine and exponential integrals, *Math. Gazette* **99** (2015), 276–289.
- [Jam2] G. J. O. Jameson, Using double integrals to solve single integrals, *Math. Gazette* **100** (2016), 257–265.
- [Jam3] G. J. O. Jameson, The incomplete gamma functions, *Math. Gazette* **100** (2016), 298–306.
- [Lag] Jeffrey C. Lagarias, Euler’s constant: Euler’s work and modern developments, *Bull. American Math. Soc.* **50** (2013), 527–628.
- [Lo] Nick Lord, Intriguing integrals: an Euler-inspired odyssey, *Math. Gazette* **91** (2007) 415–427.
- [Sco] J. A. Scott, Evaluating the Euler constant, *Math. Gazette* **94** (2010), 304–306.
- [WW] E. T. Whittaker and J. M. Watson, *A Course of Modern Analysis*, 4th ed., Cambridge Univ. Press (2004).

updated 8 November 2017